By using this website, you agree that EDP Sciences may store web audience measurement

cookies and, on some pages, cookies from social networks. More information and setup

Sciences Journals Books

Conferences



MATEC Web of Conferences

All issues Series Forthcoming About





All issues > Volume 197 (2018)

Previous issue

Free Access to the whole issue

Table of Contents

Next issue >

MATEC Web of Conferences

Volume 197 (2018)

The 3rd Annual Applied Science and Engineering Conference (AASEC 2018)

Bandung, Indonesia, April 18, 2018 Ade Gafar Abdullah and Asep Bayu Dani Nandiyanto (Eds.)

Export the citation of the selected articles Export Select all

Open Access

Statement of Peer review Published online: 12 September 2018 PDF (34.4 KB)

Open Access

About the conference Published online: 14 September 2018 PDF (10.7 MB)

OK

Industry Engineering

~ Architecture

Information Engineering

Computer and Communication Engineering

By using this website, you agree that EDP Sciences may store web audience measurement

- cookies and, on some pages, cookies from social networks. More information and setup
 Physics
 Computer Science
 Material Science
 Material Science
 Environmental Engineering
- Chemistry
- ✓ Biology
- Pharmacology
- ✓ Mechanical Engineering
- Chemical Engineering

- Mathematics

Open Access

Solving the heart curve with geogebra 01001

Ida Nuraida, Hamdan Sugilar, Rahayu Kariadinata, Juariah Juariah and Nunung Sobarningsih Published online: 12 September 2018 DOI: https://doi.org/10.1051/matecconf/201819701001 PDF (411.9 KB) References

Open Access

Representations of cyclically ordered semigroups 01002Rizky Rosjanuardi, Isnie Yusnitha and Sumanang Muhtar GozaliPublished online: 12 September 2018DOI: https://doi.org/10.1051/matecconf/201819701002PDF (240.6 KB)References

Open Access

ОK

The kumjian-pask algebras of finite finitely aligned k-graphs without cycle as modules 01003 Sumanang Gozali, Rizky Rosjanuardi and Isnie Yusnitha Published online: 12 September 2018 DOI: https://doi.org/10.1051/matecconf/201819701003 PDF (255.3 KB) References

Open Access

The behaviour of measles transmission in three different populations 01004 Budi Priyo Prawoto, Dimas Avian Maulana and Yuliani Puji Astuti Published online: 12 September 2018 DOI: https://doi.org/10.1051/matecconf/201819701004 PDF (298.4 KB) References By using this website, you agree that EDP Sciences may store web audience measurement

cookies and, on some pages, cookies from social networks. More information and setup PDF (341.5 KB) References

- Environmental Engineering

Open Access

Soil quality in corn cultivation using bamboo biochar, compost, and phonska 13001 Yohanes P Situmeang Published online: 12 September 2018 DOI: https://doi.org/10.1051/matecconf/201819713001 PDF (199.3 KB) References

Open Access

The effectiveness of "Simantri" program in producing organic fertilizer and biourine to organic farming system in Bali, Indonesia 13002

I D N Sudita, Y T L Suariani, I N Kaca and N M Yudiastari Published online: 12 September 2018 DOI: https://doi.org/10.1051/matecconf/201819713002 PDF (198.7 KB) References

Open Access

Coastline change analysis and erosion prediction using satellite images 13003 Putu Aryastana, I Made Ardantha and Kadek Windy Candrayana Published online: 12 September 2018 DOI: https://doi.org/10.1051/matecconf/201819713003 PDF (364.0 KB) References

Open Access

Design of water hyacinth utilization as renewable electricity energy source in Tondano Lake 13004 Herry Sumual, Felly Warouw and Marlon Kamagi Published online: 12 September 2018 DOI: https://doi.org/10.1051/matecconf/201819713004 PDF (216.9 KB) References

Open Access

The application of grey water recycling at Gayanti City Apartment, South Jakarta 13005 A Muthiya, T Tazkiaturrizki and R Ratnaningsih Published online: 12 September 2018 DOI: https://doi.org/10.1051/matecconf/201819713005

OK

By using this website, you agree that EDP Sciences may store web audience measurement

cookies and, on some pages, cookies from social networks. More information and setup

Open Access

Mapping the potential of green energy to border societies of Indonesia and Timor Leste (a preliminary study) 13006 Frederikus Fios

Published online: 12 September 2018 DOI: https://doi.org/10.1051/matecconf/201819713006 PDF (170.8 KB) References

Open Access

Analysis of water quality in Maninjau Lake, West Sumatera, Indonesia using phytoplankton 13007 Muhammad Arif Salsabil, Astri Rinanti and Melati Ferianita Fachrul Published online: 12 September 2018 DOI: https://doi.org/10.1051/matecconf/201819713007 PDF (235.8 KB) References

Open Access

Potential of biomass residues from oil palm agroindustry in Indonesia 13008 Dwi Ermawati Rahayu, Dutarama Nasarani, Wahyono Hadi and Budisantoso Wrjodirjo Published online: 12 September 2018 DOI: https://doi.org/10.1051/matecconf/201819713008

PDF (395.8 KB) References

Open Access

Design of drinkable tap water system at X Apartment, Jakarta 13009Hariastuti Prameswari, Tazkiaturrizki and R RatnaningsihPublished online: 12 September 2018DOI: https://doi.org/10.1051/matecconf/201819713009PDF (213.7 KB)References

Open Access

Bioethanol production from mixed culture microalgae biomass with temperature hydrolysis variation 13010

I Gede Aditya Juliarnita, Rositayanti Hadisoebroto and Astri Rinanti

Published online: 12 September 2018

DOI: https://doi.org/10.1051/matecconf/201819713010

PDF (249.6 KB) References

Open Access

OK



Editorial board

Rachid Bennacer École Normale Supérieure, Cachan, France website

Chérifa Boukacem-Zeghmouri Université Claude Bernard Lyon 1, Villeurbanne, France website

Vladimir Buzek Slovak Academy of Sciences, Bratislava, Slovakia website

Heidi Gautschi Haute Ecole Pédagogique de Lausanne, Switzerland

Éric Lichtfouse National Research Institute for Agriculture, Food and Environment (INRAE), Aix-en-Provence, France website

Maria S. Madjarska Seoul National University, Department of Physics & Astronomy, Korea And Max Planck Institute for Solar System Research, Germany By using this website, you agree that EDP Sciences may store web audience measurement cookies and, on some pages, cookies from social networks. More information and setup

Jun Sun Tianjin University of Science and Technology, P.R. China website

Ming-Jun Zhang DGUT-CNAM Institute, Dongguan University of Technology, Guangdong Province, P.R. China website

Zhien Zhang The Ohio State University, Columbus, USA

MATEC Web of Conferences

eISSN: 2261-236X Copyright / Published by: EDP Sciences Mentions légales Contacts Privacy policy





Statement of Peer review

In submitting conference proceedings to *Web of Conferences*, the editors of the proceedings certify to the Publisher that

- 1. They adhere to its **Policy on Publishing Integrity** in order to safeguard good scientific practice in publishing.
- 2. All articles have been subjected to peer review administered by the proceedings editors.
- 3. Reviews have been conducted by expert referees, who have been requested to provide unbiased and constructive comments aimed, whenever possible, at improving the work.
- 4. Proceedings editors have taken all reasonable steps to ensure the quality of the materials they publish and their decision to accept or reject a paper for publication has been based only on the merits of the work and the relevance to the journal.

The 3rd Annual Applied Science and Engineering Conference (AASEC) 18 April 2018, Universitas Pendidikan Indonesia, Bandung, Indonesia

Proceedings editor(s): Ade Gafar Abdullah Asep Bayu Dani Nandiyanto

Bandung, 12 August 2018

Ade Gafar Abdullah



CERTIFICATE No. 25/UN40.B.17/TU/2018

This certificate is awarded to

Y P Situmeang

as Presenter of a paper entitled

Soil quality in corn cultivation using bamboo biochar, compost, and phonska

in the Annual Applied Science and Engineering Conference "Ideas for Sustainable Green Energy" Bandung, Indonesia. April 18th 2018



Vice Rector for Research, Partnership, and Business Universitas Pendidikan Indonesia



Dr. Ade Gafar Abdullah, M.Si Conference Chair

(E) Engineering Village Google









Soil quality in corn cultivation using bamboo biochar, compost, and phonska

Yohanes P Situmeang¹

¹Universitas Warmadewa, Agriculture Faculty, Denpasar-Bali, Indonesia

Abstract. This study aims to determine the status of soil quality after biochar treatment, compost, and phonska in the cornfield. Soil samples were taken from 48 experimental plots after harvesting of maize. Assessment of soil quality is done by collecting selected indicator data including soil physical and chemical properties to observe changes in soil due to land use and agricultural cultivation practices. The soil quality status in this study was determined by calculating the value of soil quality rating (SQR) based on the weighting of 11 indicators of soil quality. The results showed that soil quality before the research was bad (SQR 30) and improved to moderate (SQR 27) until a good (SQR 20) after the research of biochar, compost and NPK phonska on the cornfield. Improved soil quality in biochar formulations 10.52 t ha-1, compost, and phonska, due to improved soil physical properties such as porosity, bulk density, and soil moisture content, which has encouraged the process of exchange and chemical reactions in the soil to release nutrients for the plant. Giving of bamboo biochar, compost, and NPK phonska on dry land cultivated maize has resulted in good soil quality (SQR 20) or approaching sustainability status is very good (highly sustainable).

1 Introduction

Sustainable agriculture is an effort made to maintain the productivity and quality of the land. Soil qualities are defined as soil capacity to function in ecosystems and land use limits, to maintain biological productivity, maintain environmental quality, and plant and animal health [1]. The quality of land in agriculture, referring to the ability of the soil to maintain production. High soil quality is associated with efficient water use, nutrition and pesticides, water and air quality improvement, greenhouse gas emissions mitigation, and increased agronomic production [2]. Soil quality cannot be measured directly, but static or dynamic soil quality indicators or measurable land attributes are generally affected by land use and soil management practices [3-5].

Soil physical and chemical properties can be used as indicators of soil quality assessment and sustainability of agricultural systems. Several sets of minimum data have been proposed to quantitatively assess the sustainability of soil management practices [6]. The criteria used for the sustainability of agricultural systems are based on the critical limits of the main soil properties in relation to the threshold value beyond which productivity is greatly reduced or drastic environmental impact [2, 5]. The determination of soil quality rating (SQR) based on the sum of the value and weight of each soil quality indicator can be used in soil quality rating [2].

Currently, balanced fertilization using organic and inorganic fertilizers to maintain soil fertility is still not fully applied to agricultural cultivation systems. Some research on balanced fertilization application has been done, such as by [7-9], by using biochar, compost and NPK phonska in the dry land can increase growth and yield of corn, and can improve physical and chemical properties in the soil. However, the dynamics of changes in soil properties still need to be assessed for soil quality to determine the sustainability of land use and agricultural systems. Based on the above description, this study aims to determine the status of soil quality and the sustainability of agricultural cultivation system on corn fields applied with biochar, compost, and Phonska.

2 Materials and methods

The materials used in this research are bamboo biochar, compost, NPK phonska, and soil samples. The soil sample is taken from the dry land of Sulahan Village, District of Susut, Bangli Regency. Analysis of physical and chemical properties of soil was carried out in the laboratory of Soil Science, Agriculture Faculty, Udayana University and the laboratory of Agricultural Faculty, Warmadewa University.

This research uses factorial randomized block design with 2 factors. The first factor was the dose of biochar (D) with four levels (without biochar, 5.26 t ha⁻¹, 10.52 t ha⁻¹, and 15.78 t ha⁻¹). The second factor is the type of fertilizer (P) with four types (without fertilizer, compost, phonska, and compost+NPK phonska). The treatment was repeated three times to obtain 48 plot experiments. After harvesting the maize [8], 48 of these experimental

© The Authors, published by EDP Sciences. This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (http://creativecommons.org/licenses/by/4.0/).

Corresponding author: <u>yohanes@warmadewa.ac.id</u>

plots were taken soil samples for testing soil properties in the laboratory and soil quality assessments.

The land quality assessment is conducted by collecting selected indicator data including physical and soil chemistry data (Table 1) and the use of such data to

observe changes in soil due to land use change and management practices with reference to [2]. The limiting factor and the relative weighting of the 11 minimum data sets (MDS) indicator of soil quality are presented in Table 1.

Table 1. Relative weighting factors (RWF) and critical levels for some soil p	properties	[2,10].
---	------------	---------

Indicator		Limitatior	n dan Relative weighting factor	s (RWF)	
	None	Slight	Moderate	Severe	Extreme
	1	2	3	4	5
1. Water content (%)	>30	20-30	8-20	2-8	<2
2. Bulk Density (g.cm ⁻³)	<1.3	1.3-1.4	1.4-1.5	1.5-1.6	<1.6
3. Porosity (%)	>20	18-20	15-18	10-15	<10
4. Soil Texture	L	SiL, Si, SiCL	CL, SL	SiC, LS	S, C
5. pH	6-7	5.8-6.0 or 7.0-7.4	5.4-5.8 or 7.4-7.8	5.0-5.4 or 7.8-8.2	<5.0 or >8.2
6. C-Organic (%)	5-10	3-5	1-3	0.5-1.0	<0.5
7. N-total (%)	>0.75	0.51-0.75	0.21-0.50	0.1-0.2	< 0.1
8. P-available (ppm)	>35	26-35	16-25	10-15	<10
9. K-available (ppm)	>390	234-390	156-195	78-117	<78
10. CEC (me100g ⁻¹)	>40	25-40	17-24	5-16	<5
11. BS (%)	>70	51-70	36-50	20-35	<20

L is loam; SiL is silt loam; SiCL is silty clay loam; CL is clay loam; SL is sandy loam; SiC is silty clay;

LS is loamy sand; C is clay; S is sand, CEC is cation exchange capacity, BS is base saturation.

The minimum data sets that are the key characteristics of soil quality in the topical area, consist of (1) physical characteristic: moisture content, volume weight, porosity, and soil texture, (2) chemical identification: pH, C-organic, nutrient availability (P and K), CEC, and BS. The soil quality indicator is selected from soil properties indicating the capacity of the soil function or limiting factor for crop yield. The limiting factor or relative weighting factor (RWF) ranges from no limiting to extreme factor with weighting on a scale of 1 to 5 (Tables 1 and 2). The upper limit (weight 5) for soil properties with many limiting factors means that the soil quality is very bad and unsustainable, and the lower limit (weight 1) for soil properties that do not have limiting factors means soil quality is very good and sustainable.

Table 2. The sustainability of the agricultural system is relatedto the SQR based on 11 MDS [2].

Sustainability (soil quality)	RWF	SQR
Highly sustainable (very good)	1	<20
Sustainable (good)	2	20-25
Sustainable with high input (moderate)	3	25-30
Sustainable with another land use (bad)	4	30-40
Unsustainable (very bad)	5	>40

Soil quality status was determined by calculating the value of soil quality rating (SQR) based on the sum of the weight of the value of each indicator of soil quality. Furthermore, the SQR value associated with the sustainability of agricultural systems according to [2] compared to the soil quality status criteria based on 11 Minimum Data Sets is presented in Table 2. This SQR value ranges from 20-40, the lower the SQR value the soil properties become the limiting factor less so the better the soil quality.

3 Results

3.1 Characteristics of soil, bamboo biochar, and compost

Sulahan Village, Susut Sub-district, Bangli District was chosen as a research location because it has a large dry land potential to be developed for food crops, especially corn crops. Research location with soil type regosol humus with the altitude of place 762 m asl, while characteristic of physical and chemical of soil location of research before experiment presented in Table 3.

 Table 3. Results of soil analysis, RWF, and SQR before the experiment [7].

Treatment	Soil	RWF
WC (%)	6.19	4
Texture	9.60	3
$BD (g \text{ cm}^{-3})$	0.97	1
Porosity (%)	63.54	1
pH	6.82	1
C (%)	2.69	3
N (%)	0.19	4
P (ppm)	31.08	2
K (ppm)	31.75	5
$CEC (cmol(+) kg^{-1})$	16.10	4
BS (%)	55.89	2
SOR		30

The result of soil physical and chemical properties analysis in Table 3 shows that the research field has bad soil quality with SQR value of 30. Poor soil quality status can be improved through the provision of organic materials such as biochar and compost or inorganic such as phonska. The results of characteristic analysis of biochar and compost are presented in Table 4.

Analysis Type	Biochar	Compost
pH	7.48	7.35
EC (mmhos cm ⁻¹)	0.77	10.92
C (%)	3.08	15.51
N (%)	0.06	0.82
C/N	51.33	18.91
P (ppm)	451.78	650.14
K (ppm)	36.07	23.26
Ca (mg kg ⁻¹)	32.20	9.28
Mg (mg kg ⁻¹)	15.88	15.16
Na (mg kg ⁻¹)	14.62	15.26
WC (%)	5.48	22.47

Table 4. Characteristics of biochar and compost [7].

Table 4 shows that biochar has a pH of H2O, C/N, K, Ca, and Mg relatively higher than in compost, otherwise compost has EC, C-organic, N-total, P-available, Na, and relative water content higher than biochar. Both types of

https://doi.org/10.1051/matecconf/201819713001

organic materials, biochar and compost have different characteristics, but these two materials have the same purpose and complement each other in improving soil properties in dry land, ie increasing the ability of soil in storing water and nutrients, increasing porosity and decreasing the weight of volume soil, increase Corganic, N, P, K, CEC, and BS in the soil.

3.2 Soil quality rating (SQR)

The soil quality rating is calculated based on the total number of selected soil quality indicators as the minimum data set [2]. A minimum data set consisting of 11 selected soil physical and chemical properties determines the status of soil quality. The results of the SQR measurements or the soil quality ratings from each of the biochar dose combination treatments and with the type of fertilizer (DP) on the soil after the study can be seen in Table 5.

Table 5. Soil analysis, relative weighting factors	s (RWF), and soil	l quality rating	(SQR) on b	iochar dosa	ge treatment :	and type of
	fertilizer after th	e experiment.				

	WC	Texture	BD	Porosity	pН	С	Ν	Р	K	CEC	BS	
Treatment	%		g cm ⁻³	%		%	%	ppm	ppm	cmol(+) kg ⁻¹	%	SQR
D ₀ P ₀	7.04	11.68	0.99	62.66	6.62	3.35	0.13	30.49	444.07	15.46	23.83	
	(4)	(3)	(1)	(1)	(1)	(2)	(4)	(2)	(1)	(4)	(4)	27
D_0P_1	8.38	9.08	0.92	65.23	6.63	3.64	0.17	40.65	746.98	17.15	35.96	
	(3)	(3)	(1)	(1)	(1)	(2)	(4)	(1)	(1)	(3)	(4)	24
D ₀ P ₂	8.77	15.72	0.97	63.58	6.66	3.84	0.19	35.98	331.41	17.18	49.76	
	(3)	(3)	(1)	(1)	(1)	(2)	(4)	(1)	(2)	(3)	(3)	24
D ₀ P ₃	9.16	13.00	0.97	63.38	6.55	3.71	0.21	49.34	605.67	13.91	61.64	
	(3)	(3)	(1)	(1)	(1)	(2)	(3)	(1)	(1)	(4)	(2)	22
D_1P_0	8.64	8.72	0.91	65.56	6.72	3.82	0.16	32.76	674.56	16.17	37.14	
	(3)	(3)	(1)	(1)	(1)	(2)	(4)	(2)	(1)	(4)	(3)	25
D_1P_1	9.43	15.59	0.91	65.51	6.76	4.24	0.18	42.56	997.94	17.55	39.86	
	(3)	(3)	(1)	(1)	(1)	(2)	(4)	(1)	(1)	(3)	(3)	23
D_1P_2	8.96	10.39	0.95	64.07	6.77	3.87	0.18	64.70	710.76	17.48	35.99	
	(3)	(3)	(1)	(1)	(1)	(2)	(4)	(1)	(1)	(3)	(3)	23
D1P3	9.36	9.08	0.93	64.99	6.63	4.08	0.18	37.09	934.42	16.87	65.12	
	(3)	(3)	(1)	(1)	(1)	(2)	(4)	(1)	(1)	(4)	(2)	23
D_2P_0	9.22	11.68	0.90	66.07	6.74	3.42	0.19	26.00	947.19	16.19	29.69	
	(3)	(3)	(1)	(1)	(1)	(2)	(4)	(3)	(1)	(4)	(4)	26
D_2P_1	9.71	11.69	0.89	66.30	6.68	3.84	0.17	68.06	1010.59	18.67	63.69	
	(3)	(3)	(1)	(1)	(1)	(2)	(4)	(1)	(1)	(3)	(2)	22
D_2P_2	9.87	9.09	0.89	66.45	6.75	3.83	0.20	38.99	695.08	17.57	47.69	
	(3)	(3)	(1)	(1)	(1)	(2)	(3)	(1)	(1)	(3)	(3)	22
D_2P_3	10.41	11.29	0.89	66.57	6.79	3.80	0.20	63.38	1247.02	17.18	71.73	
	(3)	(3)	(1)	(1)	(1)	(2)	(3)	(1)	(1)	(3)	(1)	20
D ₃ P ₀	9.61	12.98	0.88	66.84	6.56	3.41	0.19	38.18	1026.34	16.57	31.87	
	(3)	(3)	(1)	(1)	(1)	(2)	(4)	(1)	(1)	(4)	(4)	25
D_3P_1	10.40	12.97	0.87	67.31	6.62	3.68	0.18	40.75	1046.57	22.89	37.74	
	(3)	(3)	(1)	(1)	(1)	(2)	(4)	(1)	(1)	(3)	(3)	23
D ₃ P ₂	9.03	12.98	0.86	67.48	6.79	3.73	0.25	42.30	1018.34	18.04	46.49	
	(3)	(3)	(1)	(1)	(1)	(2)	(3)	(1)	(1)	(3)	(3)	22
D ₃ P ₃	9.01	10.35	0.87	67.29	6.74	3.77	0.24	41.39	1100.93	19.21	22.07	
	(3)	(3)	(1)	(1)	(1)	(2)	(3)	(1)	(1)	(3)	(4)	23

 D_0 (without biochar), D_1 (5.26 t ha⁻¹), D_2 (10.52 t ha⁻¹), D_3 (15.78 t ha⁻¹), P_0 (without fertilizer), P_1 (compost 20.22 t ha⁻¹), P_2 (phonska 313.37 kg ha⁻¹), dan P_3 (compost+phonska), WC is water content, BD is bulk density, CEC is cation exchange capacity, BS is base saturation, SQR: <20 = very good, 20-25 = good, 25-30 = moderate, 30-40 = bad, >40 = very bad.

4 Discussion

Assessment of soil quality before the research obtained SQR value 30 (Table 3) with bad status. This poor status after the trial can be improved to moderate to good status with a range of SQR values of 20-27 at various biochar doses and fertilizer types (Table 5). In the treatment of D_2P_3 (Table 5), the value of SQR 20 (good) or close to the status of sustainability is very good (highly sustainable), whereas in treatment D_0P_0 obtained SQR 27 (moderate). The lower the SQR score the better the sustainability index for corn farming in the dry land, the higher the SQR value the worse the sustainability index will be.

Improved soil quality from moderate to D_0P_0 treatment to good on D_2P_3 treatment is due to biochar and compost according to its characteristics (Table 4) is able to improve soil physical and chemical properties that provide the balance and nutrient adequacy needed by corn crops. Improved soil quality due to biochar 10.52 t ha⁻¹, compost and phonska (D_2P_3) fertilizers into the soil, starting from improving the quality of soil physical properties such as porosity, bulk density, and soil moisture content, which encourages the increasing process of exchange and reaction chemical in the soil to provide nutrients for plants.

Improved soil moisture due to biochar and compost treatment resulted in soil porosity ($r = 0.65^*$) and K-

available ($r = 0.67^*$) was positively and significantly correlated, but the soil bulk density ($r = -0.65^*$) was negatively correlated. Improved soil porosity due to biochar and compost treatment also caused K-available $(r = 0.82^{**})$ to be positively and highly correlated and CEC ($r = 0.63^*$) was positively and tangibly correlated, but otherwise the bulk density was negatively correlated with K- is available ($r = -0.82^{**}$) and has a negative and real correlation with CEC ($r = -0.63^*$) (Table 6). This is in line with [9, 11], that soil quality improvement due to biochar treatment is supported by the physical characteristics of biochar morphology in SEM 2000x enlargement with the surface area and micropore structure scattered on the biochar surface. These porous biochar pores lead to improved aeration and drainage systems, as well as increased soil ability to absorb ions and water in the soil. Furthermore [12] adds that bamboo biochar has a very microporous structure, with adsorption efficiency about ten times higher than traditional wood biochar. Biochar bamboo can improve aggregation and hold the capacity of groundwater, pH, and CEC soil and increase soil biological activity [13, 14]. Biochar can improve soil carbon capability, maintain soil ecosystem balance, and improve soil fertility, and can act as fertilizer, promote growth and crop yields by providing and maintaining nutrients in the soil [15, 16].

Table 6. The coefficient of correlation between soil properties.

	Water content	Texture	BD	Porosity	pН	С	Ν	Р	K	CEC
Texture	0.10									
BD	-0.65*	0.11								
Porosity	0.65*	-0.11	-1.00							
pН	0.18	-0.11	-0.36	0.36						
С	0.34	0.07	-0.03	0.03	0.37					
Ν	0.40	0.13	-0.55	0.55	0.35	0.16				
Р	0.41	-0.01	-0.07	0.07	0.23	0.34	0.10			
K	0.67*	-0.06	-0.82**	0.82**	0.32	0.15	0.44	0.32		
CEC	0.49	0.08	-0.63*	0.63*	0.19	0.18	0.20	0.16	0.47	
BS	0.50	0.03	0.03	-0.03	0.00	0.47	0.12	0.54	0.16	-0.12
			r (0.	.01, 10, 1) = 0.70	08				

5 Conclusions

The quality of the soil prior to the study had a bad status (SQR 30) and after the biochar, compost and NPK phonska studies had improved to moderate status (SQR 27) to good (SQR 20). Giving bamboo biochar 10.52 t ha⁻¹ along with compost 20.22 t ha⁻¹ and NPK phonska 313.37 kg ha⁻¹ has yielded good soil quality (SQR 20) or approaching highly sustainable status dry land cultivated corn crops.

Thanks to the Kemenristekdikti DRPM who helped fund the research in 2016. Thanks also to the students and laboratory analysts who have helped this research.

References

1. J W Doran and T B Parkin, Defining and assessing

soil quality p. 3-21 In: J.W. Doran et al., (ed.) Defining Soil Quality for a Sustainable Environment. SSSA Spec. Publ. No. **35**, Soil Sci. Soc. Am., Inc. and Am. Soc. Agron., Inc., Madison. (1994)

- R Lal No. 631.4 L193m. 85p (Washington US: Soil Management Support Services, 1994)
- C A Seybold, M J Mausbach, D L Karlen, and H H Rogers, Quantification of soil quality p 387-404 In: R. Lal, et al (Eds.) Soil Processes and the Carbon Cycle. (CRC Press, Boca Raton, 1998)
- M Sanchez-Maranon, M Soriano, G Delgado, and R Delgado Soil Science Society of America Journal 66 3 p948-958 (2002)
- 5. M K Shukla, R Lal, and M Ebinger, Soil Tillage

Research 87 2 p194-204 (2006)

- 6. W E Larson and F J Pierce, The dynamics of soil quality as a measure of sustainable management p37-51, (1994)
- Y P Situmeang, I M Adnyana, I N N Subadiyasa, and I N Merit, Int. J. on Advanced Science, Engineering and Information Technology 5 6 p433-439 (2015)
- 8. Y P Situmeang *Doctoral dissertation* p174 (Universitas Udayana, 2017)
- Y P Situmeang, Int. Research J of Engineering, IT & Scientific Research (IRJEIS) 3 3 p38-48 (2017)
- Balai Penelitian Tanah. Badan Penelitian Pengembangan Pertanian p136 (Departemen Pertanian. Bogor, 2009)
- 11. Y P Situmeang J of Biological and Chemical Research **34** 2 p704-712 (2017)
- L Hua, W Wu, Y Liu, M B McBride, and Y Chen Environmental Science and Pollution Research 16 p1–9 (2009)
- K Y Chan, B L Van-Zwieten, I Meszaros, D Downie, and S Joseph, Australian J of Soil Research 46 p437- 444 (2007)
- A Masulili, W H Utomo, and Syekhfani J of Agriculture Science 3 p25-33 (2010)
- J Major, C Steiner, A Ditommaso, N P Falcao, and J Lehmann, Weed Biol Manag 5 p69-76 (2005)
- C Steiner, W Teixeira, J Lehmann, T Nehls, J De-Macdo, W Blum, and W Zech, *Plant and Soil* 291 p275-290 (Springer Netherlands, 2007)