

# Utilization of fermented cassava skin ration on carcass weight and percentage of broiler carcass

*by Ni Made Ayu Gemuh Rasa Astiti*

---

**Submission date:** 19-Apr-2021 09:43AM (UTC+0700)

**Submission ID:** 1563035217

**File name:** n\_ration\_on\_carcass\_weight\_and\_percentage\_of\_broiler\_carcass.pdf (2.22M)

**Word count:** 8447

**Character count:** 46070

PAPER • OPEN ACCESS

## Preface

To cite this article: 2021 *J. Phys.: Conf. Ser.* **1869** 011001

View the [article online](#) for updates and enhancements.



The Electrochemical Society  
Advancing solid state & electrochemical science & technology

**240th ECS Meeting** ORLANDO, FL

Orange County Convention Center Oct 10-14, 2021

Abstract submission deadline extended: April 23rd

SUBMIT NOW

## Preface

This present volume contains the proceeding of the 2<sup>nd</sup> Annual Conference of Science & Technology (Ancoset) 2020. In this year, the 2<sup>nd</sup> Ancoset was virtually delivered through online platform due to Covid-19 Pandemic on November 28, 2020. This conference was organized by Universitas PGRI Kanjuruhan Malang and co-organized by Rumah Publikasi Indonesia.

Taking the theme of *The Role of Science and Technology Application on Building Community Resilience during New Normal*, we provided intellectual forum for students, researchers, lecturers, and practitioners. There are 221 papers collected which propose an insightful thought on the frontier knowledge about science and technology and its application. This conference opens an opportunity for scientific cooperation as well as international collaboration as it attended by almost two hundreds participants both across Indonesia and Asia.

We really appreciate all support given for the success of the 2<sup>nd</sup> Ancoset 2020 and thus would like to express our gratitude to everyone who has taken part in this event:

- The scientific reviewers for helping authors come up with a good quality paper,
- The distinguished keynote speakers: **Assoc. Prof. Sudi Dul Aji, M.Pd** (Universitas PGRI Kanjuruhan Malang), **Assoc. Prof. Dr. Massudi Mahmuddin** (Universiti Utara Malaysia), **Prof. Ida Hamidah, M.Si** (Universitas Pendidikan Indonesia), **Prof. Dr. Ade Gafar Abdullah, M.Si** (Universitas Pendidikan Indonesia),
- Co-Host universities: Universitas Al-Ghifari, Universitas Hamzanwadi, Universitas PGRI Semarang, Institut Teknologi Nasional Malang, Universitas Ma Chung, Universitas Bhinneka PGRI, Universitas PGRI Adi Buana Surabaya, IKIP PGRI Jember,
- All presenters and participants, and
- The committee members for the commitment, effort, and hard work in accomplishing the conference and proceeding.

We wish you all a fruitful gathering and see you on the next conference.

The Editors,

**Prof. Dr. Ade Gafar Abdullah, M.Si.**  
**Asst. Prof. Ayu Liskinasih, M.Pd.**  
**Asst. Prof. Muhammad Nur Hudha, M.Pd.**

**LIST OF COMMITTEE****Conference Chair:**

Asst. Prof. Ayu Liskinasih, M.Pd.

**Co-Conference Chair:**

Asst. Prof. Muhammad Nur Hudha, M.Pd.

**Advisory Boards:**

Prof. Dr. Duran Corebima Aloysius, M.Pd. - Universitas PGRI Kanjuruhan Malang

Prof. Suhadi Ibnu, Ph.D. - Universitas PGRI Kanjuruhan Malang

Prof. Laurens Kaluge, M.A., Ph.D. - Universitas PGRI Kanjuruhan Malang

Prof. Dr. Soedjijono, M.Hum. - Universitas PGRI Kanjuruhan Malang

Prof. Dr. Azlinda Azman - Universiti Sains Malaysia

Prof. Reevany bin Bustami, Ph.D. - Universiti Sains Malaysia

Prof. Dr. Mustafa bin Mamat - UNISZA

Assoc Prof. Helen Creese, Ph.D. - University of Queensland

Dr. Sam Wane - Harper Adams University

Assoc Prof. Pieter Sahertian, M.Si. - Universitas PGRI Kanjuruhan Malang

Assoc Prof. Dr. Sudi Dul Aji, M.Si. - Universitas PGRI Kanjuruhan Malang

Assoc. Prof. Dr. Muhdi, S.H., M.Hum. - Universitas PGRI Semarang

Assoc Prof. Dr. Ir. Hj. Sitti Rohmi Djalilah, M.Pd. - Universitas Hamzanwadi

Assoc. Prof. Dr. Murphin Joshua Sembiring, M.Si. - Universitas Ma Chung

Assoc Prof. Dr. H. Didin Muhafidin, M.Si. - Universitas AL-Ghifari

Assoc Prof. Dr. Ir. Kustamar, M.T. - Institut Teknologi Nasional Malang

Assoc Prof. Dr. Imam Sujono, S.Pd., M.M. - Universitas Bhinneka PGRI

Assoc Prof. Dr. M. Subandowo, M.S. - Universitas PGRI Adi Buana

Prof. Dr. H. M. Rudy Sumiharsono, M.M. - IKIP PGRI Jember

**Scientific Committee:**

Prof. Dr. Ade Gafar Abdullah, M.Si. - Universitas Pendidikan Indonesia

Assoc. Prof. Dr. Isma Widiaty, M.Pd. - Universitas Pendidikan Indonesia

Prof. Dr. Anna Permanasari, M.Si. - Universitas Pendidikan Indonesia

Asst. Prof. Muhammad Nur Hudha, M.Pd. - Universitas PGRI Kanjuruhan Malang

Asst. Prof. Ayu Liskinasih, M.Pd. - Universitas PGRI Kanjuruhan Malang

**Organizing Committee:****Secretary and Moderator:**

Asst. Prof. Riza Weganofa, M.Pd. - Universitas PGRI Kanjuruhan Malang

**Treasurer:**

Asst. Prof. Ati Retna Sari, S.E., Ak., M.SA. - Universitas PGRI Kanjuruhan Malang

**Program Committee and Information Technology:**

- Asst. Prof. Arief Rahman Hakim, M.Pd. - Universitas PGRI Kanjuruhan Malang
- Asst. Prof. Hestiningtyas Yuli Pratiwi, M.Pd. - Universitas PGRI Kanjuruhan Malang
- Asst. Prof. Tri Ida Wahyu Kustyorini, S.Pt., M.P. - Universitas PGRI Kanjuruhan Malang
- Imam Ariffudin, M.Pd. - Universitas PGRI Kanjuruhan Malang
- Ajeng Intan Nur Rahmawati, M.Pd. - Universitas PGRI Kanjuruhan Malang
- Akhmad Zaini, S.Kom., M.T. - Universitas PGRI Kanjuruhan Malang
- Heri Santoso, M.Kom. - Universitas PGRI Kanjuruhan Malang



## VIRTUAL CONFERENCE DOCUMENTATION



ANCOSET 2020 ~ 2nd Annual Conference of Science and Technology








The screenshot shows a Zoom presentation slide. The slide has a dark blue background with a world map and a rocket. The title is "Personalized Learning: A Humble Proposal of an Integrated Big Data, AI, IoT, and Security". Below the title, it says "Presentation Annual Conference of Science and Technology (Ancoset) 2020". The presenter's name is "Assoc. Prof. Dr. Massudi Mahmuddin" and his affiliation is "Universiti Utara Malaysia". The slide also features the logos of "ancoset 2020" and "UUM Universiti Utara Malaysia". At the bottom, there is a banner that reads "THE ROLE OF SCIENCE & TECHNOLOGY APPLICATION OF BUILDING COMMUNITY RESILIENCE DURING NEW NORMAL". On the right side of the slide, there is a vertical list of participants and rooms, including "Risa Wigenita, M...", "Universitas Ka...", "ROOM 7, EL...", "ROOM 12, RISKI...", "Masduki mahmu...", and "Room 11, Ye...".

The screenshot shows a Zoom presentation slide. The slide has a light yellow background with a pencil and a notepad. The title is "Biomass-based supercapacitor for electrical energy storage system". Below the title, it says "IDA HAMIDAH" and "UNIVERSITAS PENDIDIKAN INDONESIA". The presenter's name is "Prof. Dr. Ida Hamidah, M.Si" and his affiliation is "Universitas Pendidikan Indonesia". The slide also features the logos of "ancoset 2020" and "UNIKAMA UNIVERSITAS KANJURUHUN MALANG". At the bottom, there is a banner that reads "THE ROLE OF SCIENCE & TECHNOLOGY APPLICATION OF BUILDING COMMUNITY RESILIENCE DURING NEW NORMAL". On the right side of the slide, there is a vertical list of participants and rooms, including "Risa Wigenita, M...", "Universitas Ka...", "ROOM 8, ERNA...", "Masduki mahm...", "ROOM 18, SRI R...", and "Lobby".





# Table of contents

## Volume 1869

2021

[Previous issue](#)[Next issue](#)

**1**

**2nd Annual Conference of Science and Technology (ANCOSET 2020), 28 November 2020, Malang, Indonesia**

Accepted papers received: 19 March 2021

Published online: 15 April 2021

[Open all abstracts, in this issue](#)

### **Preface**

011001

The following article is Open access

[Preface](#)

[Open abstract, Preface](#) [View article, Preface](#) [PDF, Preface](#)

011002

The following article is Open access

[Peer review declaration](#)

[Open abstract, Peer review declaration](#) [View article, Peer review declaration](#) [PDF, Peer review declaration](#)

### **Chemistry**

012001

The following article is Open access

[Utilization of chitosan clam bloodshells as a coagulant for processing electroplating waste](#)

C D Wulandari, H Setyobudiarso and M Koteldae

[Open abstract, Utilization of chitosan clam bloodshells as a coagulant for processing electroplating waste](#) [View article, Utilization of chitosan clam bloodshells as a coagulant for processing electroplating waste](#) [PDF, Utilization of chitosan clam bloodshells as a coagulant for processing electroplating waste](#)

012002

The following article is Open access

[Comparison of phenolic, flavonoid, and tannin contents from ethanol extract of Kratom stem \(\*Mitragyna speciosa\* Korth.\) and senggani flower \(\*Melastoma malabathrium\* L.\)](#)

H N Hanifah, G Hadisoebroto and L Dewi

[Open abstract, Comparison of phenolic, flavonoid, and tannin contents from ethanol extract of Kratom stem \(\*Mitragyna speciosa\* Korth.\) and senggani flower \(\*Melastoma malabathrium\* L.\)](#)  
[View article, Comparison of phenolic, flavonoid, and tannin contents from ethanol extract of Kratom stem \(\*Mitragyna speciosa\* Korth.\) and senggani flower \(\*Melastoma malabathrium\* L.\)](#)  
[PDF, Comparison of phenolic, flavonoid, and tannin contents from ethanol extract of Kratom stem \(\*Mitragyna speciosa\* Korth.\) and senggani flower \(\*Melastoma malabathrium\* L.\)](#)

012003

The following article is Open access

[Clove leaf ethanol extract \(\*Syzygium aromaticum\* L. Merr. and Perr\) is formulated as antiseptic liquid soap](#)

S Maryam, K Hartono and I E Widiyawati

[Open abstract, Clove leaf ethanol extract \(\*Syzygium aromaticum\* L. Merr. and Perr\) is formulated as antiseptic liquid soap](#)  
[View article, Clove leaf ethanol extract \(\*Syzygium aromaticum\* L. Merr. and Perr\) is formulated as antiseptic liquid soap](#)  
[PDF, Clove leaf ethanol extract \(\*Syzygium aromaticum\* L. Merr. and Perr\) is formulated as antiseptic liquid soap](#)

012004

The following article is Open access

[Mercury content of \*Sardinella lemuru\* caught in East Java and Bali waters](#)

A Sartimbul, J A Amandani, D Yona and M A Z Fuad

[Open abstract, Mercury content of \*Sardinella lemuru\* caught in East Java and Bali waters](#)  
[View article, Mercury content of \*Sardinella lemuru\* caught in East Java and Bali waters](#)  
[PDF, Mercury content of \*Sardinella lemuru\* caught in East Java and Bali waters](#)

012005

The following article is Open access

[The effect of different storage times at 5°C on the quality of yogurt with the addition of local taro starch \(\*Colocasia esculenta\*\) as stabilizer](#)

A T N Krisnaningsih, D Rosyidi, L E Radiati, P Purwadi, D P P Hadiani and R L Wae

[Open abstract, The effect of different storage times at 5°C on the quality of yogurt with the addition of local taro starch \(\*Colocasia esculenta\*\) as stabilizer](#)  
[View article, The effect of different storage times at 5°C on the quality of yogurt with the addition of local taro starch \(\*Colocasia esculenta\*\) as stabilizer](#)  
[PDF, The effect of different storage times at 5°C on the quality of yogurt with the addition of local taro starch \(\*Colocasia esculenta\*\) as stabilizer](#)

012006

The following article is Open access

[Raman spectra of polyethylene glycol/cellulose acetate butyrate biopolymer blend](#)

D Marlina, M Novita, M T Anwar, H Kusumo and H Sato

[Open abstract, Raman spectra of polyethylene glycol/cellulose acetate butyrate biopolymer blend](#)  
[View article, Raman spectra of polyethylene glycol/cellulose acetate butyrate biopolymer blend](#)  
[PDF, Raman spectra of polyethylene glycol/cellulose acetate butyrate biopolymer blend](#)  
012007

The following article is Open access

[Targeted discovery of glycoside hydrolases from enzymes reservoir of digestive gland \*Achatina fulica\* through functional metagenomic strategies](#)

M Kurniawati, A Yunus, H D Ayu, S Subandi and S Suharti

[Open abstract, Targeted discovery of glycoside hydrolases from enzymes reservoir of digestive gland \*Achatina fulica\* through functional metagenomic strategies](#) [View article, Targeted discovery of glycoside hydrolases from enzymes reservoir of digestive gland \*Achatina fulica\* through functional metagenomic strategies](#) [PDF, Targeted discovery of glycoside hydrolases from enzymes reservoir of digestive gland \*Achatina fulica\* through functional metagenomic strategies](#)  
012008

The following article is Open access

[Chemical analysis and organoleptic properties of banana blossoms \(\*Musa paradisiaca\*\) as filler for catfish \(\*Clarias sp.\*\) fish floss](#)

W W Perdana, W Widowaty, D R Sofia, Y Setiawan, D Hidayanti and A Zakaria

[Open abstract, Chemical analysis and organoleptic properties of banana blossoms \(\*Musa paradisiaca\*\) as filler for catfish \(\*Clarias sp.\*\) fish floss](#) [View article, Chemical analysis and organoleptic properties of banana blossoms \(\*Musa paradisiaca\*\) as filler for catfish \(\*Clarias sp.\*\) fish floss](#) [PDF, Chemical analysis and organoleptic properties of banana blossoms \(\*Musa paradisiaca\*\) as filler for catfish \(\*Clarias sp.\*\) fish floss](#)  
012009

The following article is Open access

[Quality evaluation for chili hybrid variety candidate cultivated on highland](#)

S T Rahayu, R Kirana and P S Levianny

[Open abstract, Quality evaluation for chili hybrid variety candidate cultivated on highland](#) [View article, Quality evaluation for chili hybrid variety candidate cultivated on highland](#) [PDF, Quality evaluation for chili hybrid variety candidate cultivated on highland](#)  
012010

The following article is Open access

[Inhibitory effect of tannin extract from \*Acacia mangium\* Willd.\) against gastrointestinal pathogenic bacteria](#)



D M Sinaga, E B Laconi and A Jayanegara

[Open abstract, Inhibitory effect of tannin extract from Acacia bark \(Acacia mangium Willd.\) against gastro-intestinal pathogenic bacteria](#) [View article, Inhibitory effect of tannin extract from Acacia bark \(Acacia mangium Willd.\) against gastro-intestinal pathogenic bacteria](#) [PDF, Inhibitory effect of tannin extract from Acacia bark \(Acacia mangium Willd.\) against gastro-intestinal pathogenic bacteria](#)

012011

The following article is Open access

[Effects of mycorrhiza biofertilizer on anthocyanin contents and yield of various red rice genotypes under aerobic irrigation systems](#)

W Wangiyana, I G P M Aryana and N W D Dulur

[Open abstract, Effects of mycorrhiza biofertilizer on anthocyanin contents and yield of various red rice genotypes under aerobic irrigation systems](#) [View article, Effects of mycorrhiza biofertilizer on anthocyanin contents and yield of various red rice genotypes under aerobic irrigation systems](#) [PDF, Effects of mycorrhiza biofertilizer on anthocyanin contents and yield of various red rice genotypes under aerobic irrigation systems](#)

012012

The following article is Open access

[Morphological characteristics, chemical and amino acids composition of flours from velvet beans tempe \(\*Mucuna pruriens\*\), an indigeneous legumes from Yogyakarta](#)

H Fitriyah, F Anwar and E Palupi

[Open abstract, Morphological characteristics, chemical and amino acids composition of flours from velvet beans tempe \(\*Mucuna pruriens\*\), an indigeneous legumes from Yogyakarta](#) [View article, Morphological characteristics, chemical and amino acids composition of flours from velvet beans tempe \(\*Mucuna pruriens\*\), an indigeneous legumes from Yogyakarta](#) [PDF, Morphological characteristics, chemical and amino acids composition of flours from velvet beans tempe \(\*Mucuna pruriens\*\), an indigeneous legumes from Yogyakarta](#)

012013

The following article is Open access

[The study of structural properties and photocatalytic activity of ZnO prepared by ultrasonic assisted precipitation method](#)

S Sanusi, S Setiadji, N D Hakim, S H R A Aziz, A Sawitri, H Aliah, E P Hadisantoso, A L Ivansyah and D G Syarif

[Open abstract, The study of structural properties and photocatalytic activity of ZnO prepared by ultrasonic assisted precipitation method](#) [View article, The study of structural properties and photocatalytic activity of ZnO prepared by ultrasonic assisted precipitation method](#) [PDF, The study of structural properties and photocatalytic activity of ZnO prepared by ultrasonic assisted precipitation method](#)

012014

The following article is Open access

[Biodiversity of prokaryotic microorganisms in whey tofu from Lembang, Sumedang and Garut West Java](#)

A Junitasari, F M Warganegara and M P Widhiastuty

[Open abstract, Biodiversity of prokaryotic microorganisms in whey tofu from Lembang, Sumedang and Garut West Java](#) [View article, Biodiversity of prokaryotic microorganisms in whey tofu from Lembang, Sumedang and Garut West Java](#) [PDF, Biodiversity of prokaryotic microorganisms in whey tofu from Lembang, Sumedang and Garut West Java](#)

012015

The following article is Open access

[Toxicity of essential oils from orange \(\*Citrus sinensis\* L. Obbeck\) and lemongrass \(\*Cymbopogon nardus\* L. Rendle\) on \*Aedes aegypti\* a vector of Dengue Hemorrhagic Fever \(DHF\)](#)

N Kurniasih, W Nuryadin, M N Harahap, A Supriadin and I Kinasih

[Open abstract, Toxicity of essential oils from orange \(\*Citrus sinensis\* L. Obbeck\) and lemongrass \(\*Cymbopogon nardus\* L. Rendle\) on \*Aedes aegypti\* a vector of Dengue Hemorrhagic Fever \(DHF\)](#) [View article, Toxicity of essential oils from orange \(\*Citrus sinensis\* L. Obbeck\) and lemongrass \(\*Cymbopogon nardus\* L. Rendle\) on \*Aedes aegypti\* a vector of Dengue Hemorrhagic Fever \(DHF\)](#) [PDF, Toxicity of essential oils from orange \(\*Citrus sinensis\* L. Obbeck\) and lemongrass \(\*Cymbopogon nardus\* L. Rendle\) on \*Aedes aegypti\* a vector of Dengue Hemorrhagic Fever \(DHF\)](#)

012016

The following article is Open access

[Extraction of bixin from annatto seeds with microwave](#)

E Erdawati, E Allanas and W Wesnina

[Open abstract, Extraction of bixin from annatto seeds with microwave](#) [View article, Extraction of bixin from annatto seeds with microwave](#) [PDF, Extraction of bixin from annatto seeds with microwave](#)

012017

The following article is Open access

[The effect of concentration and temperature on the activities of kenikir \(\*cosmos caudatus\*\) leaf extract as iron corrosion inhibitor in electrolyte solution of NaCl](#)

T Sudiarti, Y Yulianti and A Supriadin

[Open abstract, The effect of concentration and temperature on the activities of kenikir \(\*cosmos caudatus\*\) leaf extract as iron corrosion inhibitor in electrolyte solution of NaCl](#) [View article, The effect of concentration and temperature on the activities of kenikir \(\*cosmos caudatus\*\) leaf extract as iron corrosion inhibitor in electrolyte solution of NaCl](#) [PDF, The effect of concentration and temperature on the activities of kenikir \(\*cosmos caudatus\*\) leaf extract as iron corrosion inhibitor in electrolyte solution of NaCl](#)

012018

The following article is Open access

[Application of nanogold and nanosilver to reduce rheumatic arthritis: Case study at leprosy patients](#)

T Taufikurohmah, R Rusmini and D Soepardjo

[Open abstract, Application of nanogold and nanosilver to reduce rheumatic arthritis: Case study at leprosy patients](#) [View article, Application of nanogold and nanosilver to reduce rheumatic arthritis: Case study at leprosy patients](#) [PDF, Application of nanogold and nanosilver to reduce rheumatic arthritis: Case study at leprosy patients](#)

012019

The following article is Open access

[Tyrosinase inhibitory activity of \*Garcinia daedalanthera\* Pierre](#)

N S S Ambarwati, B Elya, Y Desmiaty and I Ahmad

[Open abstract, Tyrosinase inhibitory activity of \*Garcinia daedalanthera\* Pierre](#) [View article, Tyrosinase inhibitory activity of \*Garcinia daedalanthera\* Pierre](#) [PDF, Tyrosinase inhibitory activity of \*Garcinia daedalanthera\* Pierre](#)

012020

The following article is Open access

[Using PhET simulation to learning the concept of acid-base](#)

O Nuraida, G S Akbar, I Farida and S Rahmatullah

[Open abstract, Using PhET simulation to learning the concept of acid-base](#) [View article, Using PhET simulation to learning the concept of acid-base](#) [PDF, Using PhET simulation to learning the concept of acid-base](#)

012021

The following article is Open access

[Analysis of cyclamate sweeteners in elementary school student's drink in South Denpasar Subdistrict](#)

N M A S Singapurwa, I D A Yunantariningsih, I W Sudiarta, I P Candra and A A M Semariyani

[Open abstract, Analysis of cyclamate sweeteners in elementary school student's drink in South Denpasar Subdistrict](#) [View article, Analysis of cyclamate sweeteners in elementary school student's drink in South Denpasar Subdistrict](#) [PDF, Analysis of cyclamate sweeteners in elementary school student's drink in South Denpasar Subdistrict](#)

012022

The following article is Open access

[Investigating the potency of bioactive compounds from \*ficus religiosa\* as anti-inflammatory agent](#)

W Utami, H A Aziz, D Nasrudin, A Kusmawan, Z Anwar, M Maulana and M Daryanto

[Open abstract, Investigating the potency of bioactive compounds from ficus religiosa as anti-inflammatory agent](#) [View article, Investigating the potency of bioactive compounds from ficus religiosa as anti-inflammatory agent](#) [PDF, Investigating the potency of bioactive compounds from ficus religiosa as anti-inflammatory agent](#)

012023

The following article is Open access

[Technology development in thermochemistry materials](#)

B Wibawa, H Syakdiyah and S Kardipah

[Open abstract, Technology development in thermochemistry materials](#) [View article, Technology development in thermochemistry materials](#) [PDF, Technology development in thermochemistry materials](#)

012024

The following article is Open access

[Neuroscience update on minimally invasive decompression for surgical treatment of neural compression in lumbar spinal stenosis](#)

B Wibawa, G A Wibawa and H Syakdiyah

[Open abstract, Neuroscience update on minimally invasive decompression for surgical treatment of neural compression in lumbar spinal stenosis](#) [View article, Neuroscience update on minimally invasive decompression for surgical treatment of neural compression in lumbar spinal stenosis](#) [PDF, Neuroscience update on minimally invasive decompression for surgical treatment of neural compression in lumbar spinal stenosis](#)

012025

The following article is Open access

[Development of protein metabolism electronic module by flip PDF professional application](#)

T Susanti, F Kurniadewi and M Nurjayadi

[Open abstract, Development of protein metabolism electronic module by flip PDF professional application](#) [View article, Development of protein metabolism electronic module by flip PDF professional application](#) [PDF, Development of protein metabolism electronic module by flip PDF professional application](#)

012026

The following article is Open access

[Developing the Computer-Based Test \(CBT\) on the basic chemistry throughout a web-based application](#)

L Yunita, M Nurjayadi and H Hariwibowo

[Open abstract, Developing the Computer-Based Test \(CBT\) on the basic chemistry throughout a web-based application](#) [View article, Developing the Computer-Based Test \(CBT\) on the basic chemistry throughout a web-based application](#) [PDF, Developing the Computer-Based Test \(CBT\) on the basic chemistry throughout a web-based application](#)

012027

The following article is Open access

[Effect of addition of white turmeric extract against antibacterial properties in edible film sweet potato starch and whey protein](#)

V Amalia, F W Adipraja, A Supriadin, N N Khasanah and A Junitasari

[Open abstract, Effect of addition of white turmeric extract against antibacterial properties in edible film sweet potato starch and whey protein](#) [View article, Effect of addition of white turmeric extract against antibacterial properties in edible film sweet potato starch and whey protein](#) [PDF, Effect of addition of white turmeric extract against antibacterial properties in edible film sweet potato starch and whey protein](#)

012028

The following article is Open access

[Utilization of e-module in polymer synthetic based on green chemistry to improve student high level thinking](#)

D D Astuti, S Rahmatullah, Y Sukmawardani and C Z Subarkah

[Open abstract, Utilization of e-module in polymer synthetic based on green chemistry to improve student high level thinking](#) [View article, Utilization of e-module in polymer synthetic based on green chemistry to improve student high level thinking](#) [PDF, Utilization of e-module in polymer synthetic based on green chemistry to improve student high level thinking](#)

012029

The following article is Open access

[A new D-Loop mitochondrial nucleotide variation from individual at Naga and Kuta traditional village](#)

T D Rosahdi, R B Satiyarti, D D Nurbaeti and D Miharja

[Open abstract, A new D-Loop mitochondrial nucleotide variation from individual at Naga and Kuta traditional village](#) [View article, A new D-Loop mitochondrial nucleotide variation from individual at Naga and Kuta traditional village](#) [PDF, A new D-Loop mitochondrial nucleotide variation from individual at Naga and Kuta traditional village](#)

012030

The following article is Open access

[Developing renewable thermo-hydrothermic bioinorganic materials from bone wastes of slaughterhouses](#)

I W Karyasa

[Open abstract, Developing renewable thermo-hydrothermic bioinorganic materials from bone wastes of slaughterhouses](#) [View article, Developing renewable thermo-hydrothermic bioinorganic materials from bone wastes of slaughterhouses](#) [PDF, Developing renewable thermo-hydrothermic bioinorganic materials from bone wastes of slaughterhouses](#)

012031

The following article is Open access

[Effect of cooking time on the quality of \*nila nyat-nyat\* \(\*Oreochromis niloticus\*\)](#)

I G S Pandit, P A N K Permtananda and K Yudha

[Open abstract, Effect of cooking time on the quality of nila nyat-nyat \(\*Oreochromis niloticus\*\)](#)

[View article, Effect of cooking time on the quality of nila nyat-nyat \(\*Oreochromis niloticus\*\)](#)

[PDF, Effect of cooking time on the quality of nila nyat-nyat \(\*Oreochromis niloticus\*\)](#)

012032

The following article is Open access

[Ethanol and methanol levels of red dragon fruit wine \(\*Hylocereus costaricensis\*\) with the treatment of sugar and fermentation time](#)

I W Sudiarta, I W R Saputra, N M A S Singapurwa, I P Candra and A A M Semariyani

[Open abstract, Ethanol and methanol levels of red dragon fruit wine \(\*Hylocereus costaricensis\*\)](#)

[with the treatment of sugar and fermentation time View article, Ethanol and methanol levels of](#)

[red dragon fruit wine \(\*Hylocereus costaricensis\*\) with the treatment of sugar and fermentation](#)

[time PDF, Ethanol and methanol levels of red dragon fruit wine \(\*Hylocereus costaricensis\*\) with](#)

[the treatment of sugar and fermentation time](#)

012033

The following article is Open access

[Purification of Fim-C-\*Salmonella typhi\* recombinant protein with Ni-NTA resin as raw material for typhoid disease detection kit](#)

M Nurjayadi, T Setiyoto, S F Jinan, D Hardianto, A Sulfianti, K Agustini and H A El-Enshasy

[Open abstract, Purification of Fim-C-\*Salmonella typhi\* recombinant protein with Ni-NTA resin](#)

[as raw material for typhoid disease detection kit View article, Purification of Fim-C-\*Salmonella\*](#)

[typhi recombinant protein with Ni-NTA resin as raw material for typhoid disease detection kit](#)

[PDF, Purification of Fim-C-\*Salmonella typhi\* recombinant protein with Ni-NTA resin as raw](#)

[material for typhoid disease detection kit](#)

012034

The following article is Open access

[Variation in recombinant protein production volume of Fim-C-salmonella typhi as a raw material for typhoid detection kit at laboratory scale](#)

M Nurjayadi, S F Jinan, T Setiyoto, D Hardianto, A Sulfianti, K Agustini, D Sukmawati and H A

El-Enshasy

[Open abstract, Variation in recombinant protein production volume of Fim-C-salmonella typhi as](#)

[a raw material for typhoid detection kit at laboratory scale View article, Variation in recombinant](#)

[protein production volume of Fim-C-salmonella typhi as a raw material for typhoid detection kit](#)

[at laboratory scale PDF, Variation in recombinant protein production volume of Fim-C-](#)

[salmonella typhi as a raw material for typhoid detection kit at laboratory scale](#)

012035



The following article is Open access

[Phytochemical and antioxidant capacity test on turmeric extract \(\*Curcuma longa\*\) traditionally processed in Bali](#)

P A N K Permatananda, A A S A Aryastuti, P N Cahyawati, D P C Udiyani, D Wijaya, I G S Pandit and A A N M Wirajaya

[Open abstract, Phytochemical and antioxidant capacity test on turmeric extract \(\*Curcuma longa\*\) traditionally processed in Bali](#) [View article, Phytochemical and antioxidant capacity test on turmeric extract \(\*Curcuma longa\*\) traditionally processed in Bali](#) [PDF, Phytochemical and antioxidant capacity test on turmeric extract \(\*Curcuma longa\*\) traditionally processed in Bali](#)  
012036

The following article is Open access

[Development of mobile learning integration with scientific approaches on electrolyte solution and redox reaction](#)

U Cahyana, E Fitriani and W Utari

[Open abstract, Development of mobile learning integration with scientific approaches on electrolyte solution and redox reaction](#) [View article, Development of mobile learning integration with scientific approaches on electrolyte solution and redox reaction](#) [PDF, Development of mobile learning integration with scientific approaches on electrolyte solution and redox reaction](#)  
012037

The following article is Open access

[Development of mobile learning integration with scientific approach in stoichiometry](#)

U Cahyana, T Hadinugrahaningsih and R I Luthfianisa

[Open abstract, Development of mobile learning integration with scientific approach in stoichiometry](#) [View article, Development of mobile learning integration with scientific approach in stoichiometry](#) [PDF, Development of mobile learning integration with scientific approach in stoichiometry](#)  
012038

The following article is Open access

[Effect of spacing on growth and yield of several black soybean varieties \(\*Glycine max\* L Merrill\)](#)

K Maulana, M S Yulianti, A A N M Wirajaya, I B K Mahardika and Y P Situmeang

[Open abstract, Effect of spacing on growth and yield of several black soybean varieties \(\*Glycine max\* L Merrill\)](#) [View article, Effect of spacing on growth and yield of several black soybean varieties \(\*Glycine max\* L Merrill\)](#) [PDF, Effect of spacing on growth and yield of several black soybean varieties \(\*Glycine max\* L Merrill\)](#)  
012039

The following article is Open access

[Application of manure types in breeding several varieties of chili \(\*Capsicum frutescens\* L.\)](#)

M S Yulianti, A A N M Wirajaya, A U R Z Assegaf and Y P Situmeang

[Open abstract, Application of manure types in breeding several varieties of chili \(Capsicum frutescens L.\)](#) [View article, Application of manure types in breeding several varieties of chili \(Capsicum frutescens L.\)](#) [PDF, Application of manure types in breeding several varieties of chili \(Capsicum frutescens L.\)](#)

012040

The following article is Open access

[Chemical analysis of commercial quail laying ration with substitution of fermentation waste bean sprouts](#)

N K Mardewi, I G A D S Rejeki and N K S Rukmini

[Open abstract, Chemical analysis of commercial quail laying ration with substitution of fermentation waste bean sprouts](#) [View article, Chemical analysis of commercial quail laying ration with substitution of fermentation waste bean sprouts](#) [PDF, Chemical analysis of commercial quail laying ration with substitution of fermentation waste bean sprouts](#)

012041

The following article is Open access

[Utilization of fermented cassava skin ration on carcass weight and percentage of broiler carcass](#)

M B Mori, N M A G R Astiti and N K Mardewi

[Open abstract, Utilization of fermented cassava skin ration on carcass weight and percentage of broiler carcass](#) [View article, Utilization of fermented cassava skin ration on carcass weight and percentage of broiler carcass](#) [PDF, Utilization of fermented cassava skin ration on carcass weight and percentage of broiler carcass](#)

012042

The following article is Open access

[Application of yeasts isolated from fermented cocoa beans for biocontrol of pathogenic mold in chocolate fruit](#)

D Sukmawati, Z Arman, R Hasanah, M Balqis, A Setiawan, F Tafrijiyyah, R Sari, A Supiyani, N B Prihantini, S N A Husna *et al*

[Open abstract, Application of yeasts isolated from fermented cocoa beans for biocontrol of pathogenic mold in chocolate fruit](#) [View article, Application of yeasts isolated from fermented cocoa beans for biocontrol of pathogenic mold in chocolate fruit](#) [PDF, Application of yeasts isolated from fermented cocoa beans for biocontrol of pathogenic mold in chocolate fruit](#)

012043

The following article is Open access

[Cellulase-producing yeast isolated from fermented cocoa beans as biocontrol for pathogenic mold chocolate fruit collected from Sentul, Jawa Barat, Indonesia](#)

012204

The following article is Open access



[Assessing two-dimensional collisions second-years students using PhET simulations on momentum and impulse concept](#)

R Adimayuda, L Sari, A Ismail, I F Amalia, S Gumilar and A Samsudin

[Open abstract, Assessing two-dimensional collisions second-years students using PhET simulations on momentum and impulse concept](#) [View article, Assessing two-dimensional collisions second-years students using PhET simulations on momentum and impulse concept](#) [PDF, Assessing two-dimensional collisions second-years students using PhET simulations on momentum and impulse concept](#)

012205

The following article is Open access

[Band gap energy calculation of theaflavin and Cyanidin-3-Glucoside molecules as active material in Dye Sensitized Solar Cells using Density Functional Theory \(DFT\)](#)

G Rizqillah, I Muttaqien, Y S Perkasa, M Sanjaya, R Ramdhani, H Aliah and P Pitriana

[Open abstract, Band gap energy calculation of theaflavin and Cyanidin-3-Glucoside molecules as active material in Dye Sensitized Solar Cells using Density Functional Theory \(DFT\)](#) [View article, Band gap energy calculation of theaflavin and Cyanidin-3-Glucoside molecules as active material in Dye Sensitized Solar Cells using Density Functional Theory \(DFT\)](#) [PDF, Band gap energy calculation of theaflavin and Cyanidin-3-Glucoside molecules as active material in Dye Sensitized Solar Cells using Density Functional Theory \(DFT\)](#)

012206

The following article is Open access

[Ab-initio calculation of cubic and tetragonal phase of CsPbI<sub>3</sub> perovskite](#)

H R Shanaz, B W Nuryadin, M N Subkhi, P Pitriana and H Aliah

[Open abstract, Ab-initio calculation of cubic and tetragonal phase of CsPbI<sub>3</sub> perovskite](#) [View article, Ab-initio calculation of cubic and tetragonal phase of CsPbI<sub>3</sub> perovskite](#) [PDF](#)

PAPER • OPEN ACCESS

## Utilization of fermented cassava skin ration on carcass weight and percentage of broiler carcass

To cite this article: M B Mori *et al* 2021 *J. Phys.: Conf. Ser.* **1869** 012041

View the [article online](#) for updates and enhancements.



The Electrochemical Society  
Advancing solid state & electrochemical science & technology

**240th ECS Meeting** ORLANDO, FL

Orange County Convention Center Oct 10-14, 2021

Abstract submission deadline extended: April 23rd

SUBMIT NOW

This content was downloaded from IP address 114.122.132.16 on 18/04/2021 at 14:54

## Utilization of fermented cassava skin ration on carcass weight and percentage of broiler carcass

M B Mori, N M A G R Astiti\* and N K Mardewi

Faculty of Agriculture, Warmadewa University, Denpasar, Bali. 80235. Indonesia

\*ayugemuh@gmail.com

**Abstract.** The purpose of this study was to determine the effect of fermented cassava peel flour on carcass weights and parts of broiler carcasses. This study used 45 CP-707 broiler strains. The design used was a randomized design complete with 3 replications. This study uses fermented cassava peels mixed in ration with 5 level administration, namely: without fermented cassava skin flour (P0), fermented cassava skin flour 3% (P1), fermented cassava skin flour 6% (P2), fermented cassava skin flour 9 % (P3), and 12% fermented cassava peel flour (P4). The application of fermented cassava peel at 9% level in the 6-week broiler ration did not significantly affect all observed variables, but application at 6% level tended to increase carcass weight, breast weight, thigh weight, and wing weight. The fermentation of cassava peel flour at 12% in the ration had a significant effect in reducing carcass weights and weight of broilers at 6 weeks. The results showed that fermentation of cassava peel flour in rations up to 6% tended to increase carcass weight, chest weight, thigh weight, and swing weight, whereas at 12% had a significant effect in reducing carcass weight and weight of broilers at 6 weeks.

### 1. Introduction

Broilers are the largest provider of animal protein from livestock because of its rapid growth with a short life cycle compared to other livestock meat. The price is quite affordable, meets the tastes of the community and is easily available [1]. Broiler chickens are male or female chickens that have growth properties and weight gain that is fast and efficient in using rations [2]. Nowadays many food additives are a combination of prebiotics and probiotics as substitutes for growth promoter antibiotics. There are a number of agro-industrial wastes that can be used as alternatives to overcome high feed costs, one of which is cassava peel waste. Cassava skin waste is an agro-industrial waste from agricultural products that has a high enough potential in Indonesia, amounting to 3,327,188.6 tons/year [3] and containing 21.20% crude fiber, crude protein 4.80%, crude fat 1.22%, dry matter 32.61 %, calcium 0.36%, phosphorus 0.11%, and energy metabolism (EM) of 2960 kcal/kg [4]. Cassava peel has weaknesses which are high levels of crude fiber and HCN, so before it is used as feed material it needs to be processed first. Cassava skin with high crude fiber has a high oligosaccharide content [5], can be used as feed for Lactic Acid Bacteria (BAL), in the digestive tract, because BAL can digest high crude fiber content.

An important factor that must be maintained is the nutritional content of the ration. Costs in livestock up to 70% of all production costs [6]. The high price of raw material for the rations, some of which are still imported, needs to find alternatives or substitutes. The utilization of food crop waste and the food industry began to be seen as one solution to overcome the problem of providing food [7]. One of the



food and industrial wastes that can be utilized is cassava peel. The carbohydrate content of cassava peels is relatively high and can be used as an energy source for livestock. Cassava peel easily separated from the tubers with a thickness of 2-3 mm. The percentage of cassava peel produced ranged from 8-25% by weight of peeled tubers and contained nutritional content such as crude protein 28%, Ca 1.69%, metabolic energy 2700 Kcal/kg, and crude fiber 14.96% [8]. Cassava peel contains anti-nutrition in the form of cyanide acid (HCN) is very high. Therefore, it is necessary processing to reduce the anti-nutrients that exist in the cassava skin, one way of processing cassava skin is by fermentation [9]. Fermentation causes better nutritional value than its original ingredients, where microorganisms that are catabolic will break down complex components into simple ones so that they are easily digested. During this time the fermentation process has been widely used as an effort to increase the nutritional content of a feed ingredient, especially the protein content [10].

Fermentation is the process of breeding selected microorganisms on cassava skin media with certain conditions so that microorganisms can develop and change the chemical composition of the media into better nutritional value. Fermentation can reduce and eliminate HCN contained in feed ingredients [8]. The use of cassava skin fermentation products up to 10% does not result in a decrease in body weight gain of broiler chickens but at levels of use above 10% cause a decrease in body weight gain [11]. Application of fermented cassava skin to broilers can improve the carcasses of broiler chickens and does not affect feed quality [12]. The percentage of broiler carcasses ranges from 65-75% of the live weight at the time of slaughtering [13]. Based on the description above, it is necessary to conduct research into the provision of several levels of fermented cassava peel flour in broiler chickens so that it can provide a good influence on the carcass weight and parts of broiler chicken carcasses.

## 2. Materials and methods

This study used a completely randomized design with 5 treatments and 3 replications. The five treatments are Chicken without the provision of fermented cassava skin flour (P0), chicken has given fermented cassava skin flour as much as 3% (P1), 6% (P2), 9% (P3) and 12% (P4). Each test uses 3 broilers. The chicken used in this study was a CP-707 strain broiler chicken with an average body weight of  $513 \pm 5$  g, obtained from PT Tohpati Poultry Shop located on Jl. WR. Supratman 281, Denpasar.

The study was conducted in Banjar Kebon Kori Klod, Kesiman Village, East Denpasar District, Denpasar City Regency. The research was conducted for 6 weeks, from February 2 to March 14, 2018. Cages with a multi-level battery system were 3 plots and divided into 15 plots. Each enclosure is 60 cm long, 50 cm wide and 40 cm high, 30 cm high from the ground, each plot contains 3 chickens. Cages made of bamboo slats each cage is equipped with a place to eat made from 30 cm long pipe Farallon and as lighting at night used 15-watt incandescent bulbs. Under the cage, plywood is used to collect fallen chicken manure, cleaned every two days so that the chicken's health is maintained from the smell of ammonia produced from chicken manure.

Ration nutrition according to the standard of Scott et al., The ingredients of rations in the form of yellow corn, rice bran, fish meal, soybean meal, fermented cassava skin flour, coconut oil, and minerals are presented in Table 1 [14].

**Table 1.** The composition of the ingredients of the research ration.

Material Name	Treatment				
	P0 (0%)	P1 (3%)	P2 (6%)	P3 (9%)	P4 (12%)
Yellow Corn (%)	55	54	55	54	54
Cassava Skin (%)	15	14.5	12	11.5	9.5
Rice bran (%)	13	12	10	10	11
Fish flour ( % )	15.5	15	15.5	14	12
Soybean Meal (%)	0.5	0.5	0.5	0.5	0.5
Coconut oil	1	1	1	1	1
Total ( % )	100	100	100	100	100

Cassava skin fermentation process. Cassava skin is washed with clean water to remove dirt. After being drained clean, then sliced into small pieces, then dried in the sun to dry. Cassava peel that has been dried is ground to get cassava skin flour. Cassava skin flour steamed for 30 minutes counted when the water vapor starts to come out of the steaming surface after it is removed then cooled. After being cold the cassava skin flour weighed 3 kg is put into the bucket with 4 ml probiotics added. Stirring until evenly distributed. After evenly stored in a closed container for 14 days. After that, the fermented cassava skin is opened and aerated. The fermentation process will not only reduce or eliminate cyanide acid levels but also increase the nutritional value contained in cassava skin. The composition of feed substances in the ration is presented in Table 2.

**Table 2.** Composition of feed substances in the study ration.

Nutrition	Treatment					Standard <sup>b</sup>
	P0	P1	P2	P3	P4	
Crude Protein (%) <sup>2)</sup>	20.31	20.30	20.22	20.18	20.24	19.0 – 21
Crude Fiber (%)	4.25	4.57	4.77	5.04	5.14	5.0 – 8 <sup>2)</sup>
EM (Kcal/Kg)	3005.65	3001.9	3011.65	3011.95	3023.35	2800-3000
Fat (%)	5.82	5.51	5.43	5.35	5.27	2.0 – 7.0
Calcium (%)	0.9	0.96	1.02	1.07	1.13	0.9-1.1
Phosphor (%)	0.81	0.80	0.82	0.84	0.86	0.8-1.0

Information: 1) Based on Recommendations [14], 2) Based on Recommendations [15].

Of the 100 chickens prepared, it was first weighed to look for an average body weight of  $513 \pm 5$  gr%, then look for 45 chickens that were close to average body weight. After that 45 selected chickens were randomly placed in the experimental unit. Each experimental unit was filled with 3 chickens. A week before the chicken is put into the cage is cleaned and disinfected using Destan brand disinfectant to eradicate bacteria, viruses, pests, and fungi. When Day Old Chick (Doc) arrives given sugar water through drinking water to avoid stress, then given vita chick at a dose of 5 g, vita chick mixed with 7 liters of drinking water, the first vaccination is carried out on day 4 with the Newcastle Disease (ND) vaccine through eye drops with the aim of preventing New Castle Disease, the second and third vaccinations are carried out at 13 and 21 days with the Gumboro vaccine.

The mixing of the ration is done once a week, the material to be used is weighed according to needs. After mixing is done, the ration is put into a plastic bag that has been filled with the code according to the treatment. Feed and drinking water are given on an ad-libitum basis. Feeding is done 3 times a day, morning, afternoon and evening. The water provided comes from the local PDAM. Each ration is given half of the capacity of the feed container with the aim to avoid the rationing of the ration when the chicken is eating. The feed and drink containers are cleaned every day.

### 3. Results and discussion

To study the effect of electron velocity and magnitude of the magnetic field on the electron trajectory, the application of fermented cassava peel flour in the ratio up to level 9% shows that the results of different carcass weight were not significant, but at the level of 12%, fermented cassava peel flour produced the lowest carcass weight compared to other treatments. The application of fermented cassava peel to 12% level in the ratio showed no significant effect on the variables of breast weight, thigh weight, and back weight. But it has a significant effect in reducing carcass weight and wing weight. However, by giving 6% fermented cassava peel flour, the weight of the portion of the carcass produced in P2 is tended to be higher than other treatments. This shows that the limit of fermented cassava skin flour in broiler chicken ration is 9%. At 9% the chicken body is able to digest and decompose the ratio that is consumed properly, this is in accordance with research conducted by Supriyati and Kompiang [16]. The chest and thighs develop more dominant during growth when compared to the wings and back [17], this is because the wings and back are dominated by bone components and have less potential to produce meat. The difference in growth velocity will affect the weight distribution of body parts or carcass



components [18]. The growth of carcass components begins with bone growth, then muscle growth will decrease after reaching puberty, followed by increased fat growth [19].

**Table 3.** The average weight of carcass and carcass part of broiler fed with fermented cassava peel flour.

Variable	Treatment					SEM
	P0 (0%)	P1 (3%)	P2 (6%)	P3 (9%)	P4 (12%)	
Carcass Weight (g)	872.19 <sup>a</sup>	870.62 <sup>a</sup>	898.90 <sup>a</sup>	856.68 <sup>a</sup>	749.92 <sup>b</sup>	17.84
Chest Weight (g)	256.50 <sup>a</sup>	246.23 <sup>a</sup>	260.34 <sup>a</sup>	255.90 <sup>a</sup>	221.38 <sup>a</sup>	11.87
Thigh Weight (g)	286.37 <sup>a</sup>	283.93 <sup>a</sup>	297.87 <sup>a</sup>	294.83 <sup>a</sup>	267.30 <sup>a</sup>	7.51
Back Weight (g)	211.13 <sup>a</sup>	212.96 <sup>a</sup>	196.09 <sup>a</sup>	190.31 <sup>a</sup>	183.65 <sup>a</sup>	4.49
Wing Weight (g)	118.20 <sup>a</sup>	127.50 <sup>a</sup>	144.61 <sup>a</sup>	115.64 <sup>ab</sup>	77.59 <sup>b</sup>	6.66

Information:

Values with the same letter in the same line indicate no significant difference

SEM = Standard Error the Treatment Meant

Application of fermented cassava peel flour at a rate of 12% in broiler chicken ration showed a significantly lower effect on carcass weight. P4 treatment (12%) has the lowest carcass weight and is significantly lower than other treatments [20] which states that carcass weight is closely related to cutting weight, the high cutting weight will be followed by high carcass weight and vice versa if the weight of the slice is low then weight carcass and carcass fracture will decrease. The carcass weight produced is influenced by several factors, namely age, sex, cutting weight, body size and conformation, fat, quality and amount of ration and strain maintained [21].

The provision of fermented cassava peel flour up to 12%, can reduce the carcass weight, this is because the fermented cassava peels up to level 12% of the fiber content is too high for broilers and still has HCN content. The content of HCN has harmful antifungal agents in cattle so that it can reduce the cutting weight followed by carcass weight [5]. HCN levels that can be tolerated are no more than 50 ppm and effective processing techniques are needed, namely fermentation which can increase protein levels, digestibility and reduce HCN levels in cassava skin [22]. This is also supported by Hanifah et al., which states that HCN can suppress nitrogen retention and cause decreased digestion of amino acids which must be digested by villous intestinal villi and used for tissue growth and development [23]. Provision of 6% fermented cassava peel in the ration obtained carcass weights tended to be highest. This is caused by the fermentation process which can increase the protein content of the material and reduce the levels of crude fiber from the material that can be used by broilers to produce good carcass weights. The fermentation process can improve the digestibility of organic cassava peels as feed ingredients, fermentation techniques can eliminate HCN from feed ingredients [16].

The application of fermented cassava peel at 9% and 12% levels on the wing weight variable showed a significantly decreased effect compared to the treatments P0, P1, and P2 (Table 3). Wing weight at P4: 77.59 g/head decreased significantly compared to P0, P1, and P3. This is due to the lowest carcass weight in the P4 treatment resulting in a low wing weight. The highest wing weight was obtained at level 6% with a weight of 144.61 g/head. There is a close relationship between carcass weight and carcass parts [24], it is reinforced by Miles et al., that carcass parts are directly determined by carcass weight [25]. The decrease in carcass weight gain is also due to the higher levels of tannin and HCN in the feed because it can suppress nitrogen retention and can cause a decrease in the digestibility of amino acids [26].

#### 4. Conclusion

The provision of fermented cassava peel flour in broiler chicken ration had a significant effect on carcass weight and wing weight variables, while other variables such as breast weight and thigh weight were non-significant. The provision of fermented cassava peel flour at the 6% level tends to increase the

carcass weight and commercial portion. It is recommended that fermented cassava skin flour be at the level of 6% because it tends to increase carcass weight and commercial portion in 6 week broiler chickens.

## References

- [1] Astiti N M A G R 2018 *Pengantar Ilmu Peternakan* (Denpasar: Warmadewa University Press)
- [2] Mardewi N K, Astiti N M A G R, Rukmini N K S, Rejeki I G A D S, Tonga Y and Suwitari N K E 2017 Supplementation of moringa (*Moringa oliefera*) powder into ration in increasing the quality of broiler's meat *International research journal of engineering, IT & scientific research* **3** 3 31-44
- [3] BPS 2008 *Produksi Singkong Indonesia* (Jakarta: Badan Pusat Statistik)
- [4] Darmawan 2006 Pengaruh kulit umbi ketela pohon fermentasi terhadap tampilan kambing kacang jantan *Jurnal Ilmiah Ilmu-Ilmu Peternakan, Universitas Jambi* **9** 2 115-122
- [5] Hidayat C 2009 Peluang penggunaan kulit singkong sebagai pakan unggas *Seminar Nasional Teknologi Peternakan dan Veteriner*
- [6] Astiti N M A G R 2018 *Sapi Bali dan Pemasarannya* (Jayapangus Press Books)
- [7] Astiti N M A G R 2019 Impact of Bali Cattle Calf Marketing to the Farmers Income *The Journal of Research on the Lepidoptera* **50** 4 89-96
- [8] Supriyadi 1995 Peluang Penggunaan Kulit Singkong Sebagai Pakan Unggas *Seminar Nasional Teknologi Peternakan dan Veteriner Balai Penelitian Ternak, Bogor*
- [9] Winarno F G, Fardiaz S and Fardiaz D 1980 *Pengantar Teknologi Pangan* (Jakarta: PT. Gramedia)
- [10] Wizna, Abbas H, Rizal Y, Dharma A and Kompiang I P 2009 Improving the quality of tapioca By-Products (Onggok) as poultry feed throug fermentation by *Bacillus amyloliquefaciens* *Pakistan Journal of Nutrition* **8** 10 1636-1640
- [11] Supriyati 2003 Onggok terfermentasi dan pemanfaatannya dalam ransum ayam ras pedaging *JITV* **8** 146-150
- [12] Mirzah and Muis H 2015 Improving Nutrient Quality of Cassava Peel Waste by Fermentation Using the *Bacillus Amiloliquefaciens* *Jurnal Peternakan Indonesia* **17** 2 131-142
- [13] Mardewi N K, Rukmini N K S, Rejeki I G A D S and Astiti N M A G R 2019 The effect of cage density on the quality of broiler chicken meat *Journal of Physics: Conference Series* **1402** 3 033058
- [14] Scott M L, Nesheim M C and Young R J 1982 *Nutrition of the Chickens* Second Ed. M L Scott and Associates Ithaca (New York)
- [15] Standar Nasional Indonesia 2015 *Mutu Karkas dan Daging Ayam* (SNI)
- [16] Supriyati dan Kompiang I P 2002 Perubahan komposisi nutrien dari kulit ubi kayu terfermentasi dan pemanfaatannya sebagai bahan baku pakan ayam pedaging *Jurnal Ilmu Ternak Veteriner* **7** 3 150-154
- [17] Muryanto P, Hardjosworo S, Herman R and Setijanto H 2002 Evaluasi Karkas Hasil Persilangan Antara Ayam Kampung Jantan dengan Ayam Ras Petelur Betina *J. Anim. Prod.* **4** 2 71 – 76
- [18] Selviana E N, Suprijatna M and Mahfudz L D 2019 Pengaruh Penambahan Kulit Singkong Fermentasi dengan Bakteri Asam Laktat Sebagai Aditif Pakan Terhadap Produksi Karkas Ayam Kampung Super *Jurnal Fakultas Pertanian UNS* **3** 1 77-85
- [19] Sharifi M R, Shams-Shargh M, Dastar B dan Hassani S 2011 The effect of dietary protein levels and synbiotic on performance parameters, blood characteristics and carcass yield of Japanese quail (*Coturnix coturnix Japainica*) *Italian J. Anim. Sci.* **10** 1 478 – 488
- [20] Munira S, Nafiu L O dan Taisse A M 2016 Performans ayam kampung super pada pakan yang disubstitusi dedak padi fermentasi dengan fermentor berbeda *J. Ilmu Tropik* **3** 2 21-29
- [21] Hazim J A, Al-Mashadani H A, Al-Hayani W K, Mirza H A and Al-Hassani A S 2010 Effect of dietary supplementation with different oils on productive and reproductive performance of quail *International Journal of Poultry Science* **9** 5 429-435

- [22] Hanifah V W, Yulistiani D and Asmarasari S A A 2010 *Optimalisasi Pemanfaatan Limbah Kulit Singkong Menjadi Pakan Ternak dalam Rangka Memberdayakan Pelaku Usaha enye-enye, Prosiding Seminar Nasional Teknologi Peternakan dan Veteriner* (Bogor: Balai Penelitian Ternak)
- [23] Widodo W 2005 *Tanaman Beracun Dalam Kehidupan Ternak, Edisi Pertama* (Malang: Universitas Muhammadiyah)
- [24] Iskandar S 2006 Ayam silangan pelung – kampung : tingkat protein ransum untuk produksi daging umur 12 minggu *J. Wartazoa* **16** 2 65 -71
- [25] Miles R D, Butcher G D, Henry P R and Littell R C 2006 Effect of antibiotic growth promoters on broiler performance, intestinal growth parameters, and quantitative morphology *Poult. Sci.*, **85** 3 476-485
- [26] Astiti N M A G R, Wijaya W A and Mardewi N K 2018 Pemanfaatan Tepung Kulit Pisang Terfermentasi untuk Menekan Biaya Pakan Itik Bali Umur 2-8 Minggu in: *seminar nasional persepsi iii: Strategi dan Kebijakan Pengembangan Bisnis Peternakan Dalam Mendukung Kedaulatan Pangan Nasional* (Manado: Prosiding Universitas Sam Ratulangi)



# Utilization of fermented cassava skin ration on carcass weight and percentage of broiler carcass

## ORIGINALITY REPORT

7%

SIMILARITY INDEX

10%

INTERNET SOURCES

5%

PUBLICATIONS

3%

STUDENT PAPERS

## PRIMARY SOURCES

1

conference.unikama.ac.id

Internet Source

5%

2

Submitted to Syiah Kuala University

Student Paper

3%

Exclude quotes On

Exclude bibliography On

Exclude matches < 2%

# Utilization of fermented cassava skin ration on carcass weight and percentage of broiler carcass

## GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14

PAGE 15

PAGE 16

PAGE 17

PAGE 18

PAGE 19

