

Utilization of rabbit manure and biochar chicken manure and its effect on the growth and yield of pakchoy plants

by Yohanes Parlindungan Situmeang

Submission date: 23-Jan-2022 09:13PM (UTC+0700)

Submission ID: 1746319402

File name: 8._Mesa_2021_J._Phys._Conf._Ser._1869_012045.pdf (743.4K)

Word count: 3459

Character count: 16618

PAPER · OPEN ACCESS

Utilization of rabbit manure and biochar chicken manure and its effect on the growth and yield of pakchoy plants

To cite this article: I M Mesa *et al* 2021 *J. Phys.: Conf. Ser.* **1869** 012045

1

View the [article online](#) for updates and enhancements.

You may also like

7
- [Yields of maize \(*Zea mays* L.\) and peanut \(*Arachis hypogaea* L.\) in intercropping system treated by bokashi plus fertilizer under early growth of teak plantation in Napabalano District, Muna Regency, Indonesia](#)

L Karimuna, Halim, A Ansi *et al.*

- [Effectiveness of Biopellet Combination of Biochar, Chicken Manure and Fish Waste To Improvement of Chemical Properties of Sandy Soil and Soybean Plant Growth](#)

Sugeng Winarso, Bambang Hermiyanto, Sukron Romadhona *et al.*

6
- [Cucumber plants \(*Cucumis sativus* L.\) growth and crop yield of chicken manure fertilized with plant spacing](#)

Sri Pratiwi Arlonang, Ernitha Panjaitan and Fetrus Parsaulian Tondang



IOP ebooks™

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

Utilization of rabbit manure and biochar chicken manure and its effect on the growth and yield of pakchoy plants

I M Mesa, Y P Situmeang*, A A N M Wirajaya, I G B Udayana and M S Yuliantini

Faculty of Agriculture, Warmadewa University, Denpasar-Bali, Indonesia

*ypsitumeang63@gmail.com

Abstract. This study aims to determine the effect of rabbit manure, biochar from chicken manure, and its interactions on the growth and yield of pakchoy plants. This study uses a randomized block design (RBD) with factorial patterns (2 factors). The first factor is the dose of rabbit manure which consists of 4 levels (0, 10, 20, and 30 tons ha⁻¹). The second factor is the biochar dose of chicken manure consisting of 4 levels (0, 5, 10, and 15 tons ha⁻¹). The results showed that the treatment of biochar dose of chicken manure and the treatment of interactions between the dose of rabbit manure and the biochar dose of chicken manure had no significant effect on all observed variables. The treatment of rabbit droppings did not significantly affect all observed variables, except that the fresh weight of the roots showed a real effect. Fresh weight of economic results in the application of rabbit manure 30 tons ha⁻¹, gives a higher yield or an increase of 17.08% when compared to without rabbit manure. While the fresh weight of economic results on the application of biochar from chicken manure 30 tons ha⁻¹ gave higher yields or increased by 23.86% compared to without biochar.

1. Introduction

Pakchoy (*Brassica rapa* L.) is a group of plants from the Brassica genus used as vegetables in the form of cooking or fresh or fresh vegetables. It tastes good and distinctive and is suitable for a variety of dishes. Pakchoy plant is one of the horticultural commodities that are useful as a source of vitamins, minerals and contain fiber, such as vitamin A, vitamin B, vitamin B6, vitamin C, potassium, phosphorus, copper, magnesium, iron, and protein. The nutritional content causes pakchoy in addition to nutritious food, which is also efficacious for preventing cancer, hypertension, and heart disease. One effort to intensify agriculture that needs to be done to increase agricultural production is by fertilizing. Fertilization is done to meet the needs of nutrients in the soil so that the genetic potential of plants can be achieved to the maximum [1]. Application of cultivation techniques through the provision of organic fertilizers such as compost and biochar from animal manure in soil media can provide increased nutrient supply.

Chicken and rabbit farming on a large scale raises several problems, among others in the problem of handling cage waste, especially feces (solid waste). Cage waste in the form of livestock manure, both feces or leftover food that is scattered is the most dominant source of environmental pollution in the area of livestock. This livestock waste can be processed into organic fertilizer either as fermented compost or as a biochar soil enhancer that is useful for improving soil fertility and agricultural crop yields.



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

Biochar is black charcoal as a result of the heating process of biomass in a state of limited oxygen or no oxygen. Biochar is an organic material that has stable properties that can be used as soil amelioration in the dry land. The benefits of adding biochar to the soil include increasing plant growth, reducing methane emissions, reducing NO₂ emissions, reducing fertilizer requirements, reducing nutrient leaching, storing carbon in the long run stably, increasing soil pH, soil aggregates, soil water content, cation exchange capacity, and crop yields [2].

Utilization of biochar on a large scale is relatively new, therefore the government has an important role in providing understanding and guidance to the wider community that requires farmers to need biochar as a fixing material to increase agricultural production going forward [3]. Some compost and biochar research results in improving crop yields have been approved and published in various scientific publications. The best results of application dosage of biochar in various types of plants are biochar dose of 10 tons ha⁻¹ in maize plants [4-7], biochar dose of 10 tons ha⁻¹ in sorghum plants [8], biochar dosage of 10-15 tons ha⁻¹ in plants chili peppers [9,10], biochar 9 tons ha⁻¹ in spinach plants [11], and biochar doses 6 tons ha⁻¹ in kale plants [12] and biochar 10 tons ha⁻¹ in pakchoy plants [13], with the best results helping fertilizer 30 kg ha⁻¹ in chili plants [14,15], and compost 20 tons ha⁻¹ in pakchoy plants [13].

Chicken manure which is burned into biochar charcoal has given the best results to the chili plants [10]. Likewise, rabbit manure fermented into compost has given the best response to chili plants [14], but this may not be responded well to Pakchoy plants. Therefore this research was carried out, to obtain the best response of pakchoy plants in doses of rabbit manure and biochar from chicken manure. The hypothesis proposed in this study is that manure from rabbit manure 20 tons ha⁻¹ and biochar from chicken manure 10 tons ha⁻¹ and its interactions can increase the growth and yield of pakchoy plants.

2. Materials and methods

This research was conducted at the experimental station of the Faculty of Agriculture, Warmadewa University, Jalan Terompong, Denpasar with a height of 25 meters above sea level. This research activity began from May to June 2019. This research was a factorial experiment (two factors) using a Randomized Block Design. The first factor is the dose of rabbit compost (R) which consists of 4 levels, namely: 0 tons ha⁻¹ (R0), 10 tons ha⁻¹ (R1), 20 tons ha⁻¹ (R2), 30 tons ha⁻¹ (R3). The second factor is the biochar dose (B) which consists of 4 levels, namely: 0 tons ha⁻¹ (B0), 5 tons ha⁻¹ (B1), 10 tons ha⁻¹ (B2), and 15 tons ha⁻¹ (B3). So there are 16 treatment combinations, each repeated 3 times so there are 48 trial units. The variables observed in this study were: plant height, number of leaves, leaf area, leaf fresh weight, root fresh weight, oven-dry weight, oven-dry weight. The experimental data were analyzed according to the design used. The treatment that was significantly affected was followed by the most significant difference test (LSD) of 5%, whereas to find out the close relationship between the observed variables, correlation analysis was performed.

3. Results

Based on these results and after statistical analysis, the significance of the influence of Rabbit manure (R) and Biochar from chicken manure (B) and its interactions with the observed variables are presented in Table 1.

Table 1. Significance of the effects of rabbit manure (R) and biochar from chicken manure (B), and their interactions (RxB) on all observed variables.

No	Variable	R	B	RxB
1	Plant height (cm)	ns	ns	ns
2	Number of leaves (strand)	ns	ns	ns
3	Leaf area (cm ²)	ns	ns	ns
4	Fresh weight of economic results (g)	ns	ns	ns
5	Root fresh weight of root (g)	*	ns	ns
6	Plant oven-dry weight (g)	ns	ns	ns
7	Root oven-dry weight (g)	ns	ns	ns

Note: ns = not significant effect ($P \geq 0.05$), * = Significantly effect ($P < 0.05$)

The treatment of rabbit manure doses did not significantly affect all observed variables except that the root fresh weight had a significant effect ($P < 0.05$). However, the biochar dose treatment did not significantly affect ($P \geq 0.05$) on all observed variables. Likewise, the interaction of rabbit and biochar (RxB) manure treatments had no significant effect ($P \geq 0.05$) on all observed variables.

Statistical analysis showed that the treatment of rabbit droppings (R) showed a significant effect ($P < 0.05$), whereas the treatment of biochar dosage (B) and interactions (RxB) had no significant effect ($P \geq 0.05$) on fresh root weight. However, the fresh weight variable of economic yield per plant showed that the treatment dose of rabbit manure (R), biochar dose (B), and its interaction (RxB) had no significant effect ($P \geq 0.05$) (Table 1). The average of all plant variables observed in the care of rabbits and biochar droppings is presented in Table 2.

Table 2. Average of all variables observed by plants in the treatment of rabbit manure (R) and biochar from chicken manure (B).

Treatment	Plant height (cm)	Number of leaves (strand)	Leaf area (cm ²)	Fresh weight of economic results (g)	Root fresh weight of root (g)	Plant oven-dry weight (g)	Root oven-dry weight (g)
Rabbit manure							
0 ton ha ⁻¹ (R0)	20.87 a	9.50 a	465.96 a	51.63 a	5.16 b	2.43 a	0.65 a
10 ton ha ⁻¹ (R1)	21.80 a	9.75 a	574.39 a	53.36 a	5.08 b	2.82 a	0.73 a
20 ton ha ⁻¹ (R2)	21.53 a	9.33 a	521.94 a	56.19 a	6.07 a	2.80 a	1.00 a
30 ton ha ⁻¹ (R3)	21.38 a	10.33 a	529.14 a	60.45 a	6.41 a	2.93 a	0.95 a
LSD 5%	-	-	-	-	1.00	-	-
Biochar of chicken manure							
0 ton ha ⁻¹ (B0)	21.18 a	9.17 a	506.08 a	50.50 a	5.47 a	2.62 a	0.98 a
5 ton ha ⁻¹ (B1)	21.22 a	9.75 a	487.72 a	50.03 a	5.52 a	2.54 a	0.78 a
10 ton ha ⁻¹ (B2)	21.28 a	9.67 a	501.83 a	58.55 a	5.84 a	2.73 a	0.68 a
15 ton ha ⁻¹ (B3)	21.90 a	10.33 a	501.83 a	62.55 a	5.89 a	3.09 a	0.88 a
LSD 5%	-	-	-	-	-	-	-

From Table 2 it can be seen that the highest average fresh root weight was obtained in the treatment of rabbit manure dosages of 30 tons ha⁻¹ (R3) which is 6.41 g which is very different from the treatment of rabbit manure 10 tons ha⁻¹ (R1), which is 5.08 g and without treatment (R0) is 5.16 g, but not significantly different from the treatment of 20 tons ha⁻¹ (R2) with a value of 6.07 g. In the treatment of the highest biochar dose tends to be obtained at 15 tons ha⁻¹ (B3) which is 6.41 g not significantly different from other biochar treatment doses, while the lowest fresh root weight value without biochar (B0) is 5.47 g.

4. Discussion

The results of statistical analysis showed that the fresh weight of economic results had no significant effect on the treatment of rabbit or biochar manure (Table 1). However, the highest fresh weight of economic results tends to be obtained in the treatment of 30 tons ha⁻¹ rabbit manure (R3) at 60.45 g, an increase of 17.08% compared to without rabbit manure (R0) at 51.63 g (Table 2). The close relationship between these variables in the treatment of rabbit droppings can be seen in Table 3.

From Table 3, it can be seen that the high fresh weight of economic results is supported by the number of leaves ($r = 0.704^{**}$), fresh root weight ($r = 0.939^{**}$), oven-dry weight per plant ($r = 0.801^{**}$), and weight oven-dried roots ($r = 0.831^{**}$).

Table 3. Correlation coefficient values (r) due to the influence of rabbit manure doses.

	Plant height	Number of leaves	Leaf area	Fresh weight of Economic results	Root fresh weight	Plant oven dry weight
Number of leaves	0.128ns					
Leaf area	0.970**	0.313ns				
Fresh weight of economic results	0.258ns	0.704**	0.274ns			
Root fresh weight of root	0.090ns	0.455ns	0.036ns	0.939**		
Plant oven dry weight	0.782**	0.576*	0.789**	0.801**	0.638**	
Root oven dry weight	0.362ns	0.198ns	0.239ns	0.831**	0.919**	0.722**
	r = (0.05; 14) = 0.497			r = (0.01; 14) = 0.623		

Note: * = significantly effect ($P < 0.05$), ** = very significant effect ($P < 0.01$), ns = not significant effect ($P \geq 0.05$)

The results of this study indicate that the treatment of rabbit manure, only the fresh weight variable of the roots gives real results (Table 1). The treatment of rabbit manure 30 tons ha^{-1} (R3) gave the highest yield of fresh root weight of 6.41 g which increased by 24.22% when compared with the lowest yield in the treatment without rabbit manure (R0) which was 5.16 g.

The high fresh root weight in the treatment of rabbit manure doses was supported by the existence of a positive and real correlation on the economic fresh weight variable ($r = 0.939$ **), and the oven-dry root weight ($r = 0.919$ **) (Table 3). This can be due to the nutrient content contained in rabbit manure such as N (0.03%), P (55.56 ppm), K (59.91 ppm), and C (0.41%) is quite high. High N, P, and K nutrient content can support the supply and absorption of nutrients by roots in the soil. The nutrient content such as N, P, and K contained in rabbit feces is quite high due to a very active microbial population [16]. This is also reinforced by the statement Novizan which states that nitrogen is the main nutrient for growth, which is generally very necessary for the formation of vegetative parts of plants such as leaves, stems, and roots [17]. While phosphorus plays a role in a variety of physiological processes in plants such as photosynthesis and respiration, so does potassium plays a role in encouraging various enzymes that are essential in photosynthetic reactions

The results of this study also showed that the application of biochar from chicken manure had no significant effect ($P \geq 0.05$) on all observed variables. Nevertheless, the highest fresh weight of economic results tends to be obtained in the application of 15 tons ha^{-1} (B3) of 62.55 g or an increase of 23.86% compared to without biochar (B0) of 50.50 g (Table 2). This is due to the improved physical properties of the soil which encourage the movement of nutrients and water, so the plant roots become more active in absorbing nutrients from the soil for growth.

However, the effect of biochar in improving soil fertility takes a long time to get the best growth and crop yields. The stable and difficult to decompose biochar in the soil causes biochar residue to last for a long time in the soil. Therefore, the application of biochar is only done once for several planting seasons.

The reaction of the application of organic fertilizer requires time to be absorbed or utilized by plants because the decomposition process takes place slowly to provide nutrients for plants [18]. Various studies have been carried out showing that biochar is useful for improving soil physical properties by increasing water holding capacity and aggregate stability, improving soil weight, and reducing soil resistance due to its porous structure [19] and improving soil fertility and soil quality [20]. Biochar particles bind to very fine soil fractions of 50 μm [21] and the presence of biochar is restoring soil quality through small clusters of soil particles or aggregates compared to other organic matter [22].

5. Conclusions

The interaction between the treatment dose of rabbit manure and the biochar from chicken manure did not significantly influence all observed variables. The treatment of rabbit manure did not significantly affect all observed variables, except that the fresh weight of the roots showed a real effect. The treatment of rabbit manure 30 tons ha^{-1} gave the highest yield of fresh root weight which increased by 24.22%

from without rabbit manure. The application of rabbit manure 30 tons ha⁻¹ gives a higher fresh weight of economic yield or an increase of 17.08% when compared to without rabbit manure. Likewise, the application of biochar from chicken manure 30 tons ha⁻¹ yields higher fresh economic results or increased by 23.86% compared without biochar.

Acknowledgments

The author thanks the Head of Experimental Station of the Faculty of Agriculture, Warmadewa University and friends who have helped with this research.

References

- [1] Winaya P D 1983 *Kesuburan Tanah dan Pupuk* (Denpasar: Jurusan Ilmu Tanah, Fakultas Pertanian Universitas Udayana Denpasar)
- [2] Gani A 2009 *Pemanfaatan Arang Hayati (Biochar) Untuk Perbaikan Lahan Pertanian* (Bogor: Bahan Seminar di Puslitbangtan Bogor)
- [3] Adhi R K 2013 *Biochar Sang Pembunuh Tanah* (Balai Besar Pelatihan Binaung)
- [4] Situmeang Y P, Adnyana I M, Subadiyasa I N N and Merit I N 2015 *International Journal on Advanced Science, Engineering and Information Technology* **5** 6 433-439
- [5] Ngongo P M, Situmeang Y P and Kartini L 2018 *SEAS (Sustainable Environment Agricultural Science)* **2** 1 67-71
- [6] Situmeang Y P, Adnyana I M, Subadiyasa I N N and Merit I N 2018 *International Journal on Advanced Science, Engineering and Information Technology* **8** 5 2241-2248
- [7] Putra I G L A A A, Situmeang Y P, Yuliantini M S and Udayana I G B 2019 *Journal of Physics: Conference Series* **1402** 5 055029
- [8] Yudiastari N M, Suwitari N K E, Suariani L and Situmeang Y P 2019 *IOP Conference Series: Earth and Environmental Science* **347** 1 012057
- [9] Dapa D S, Situmeang Y P and Sudewa K A 2020 *SEAS (Sustainable Environment Agricultural Science)* **3** 2 118-123
- [10] Situmeang Y P, Sudita I D Y and Suarta M 2018 *International Journal on Advanced Science, Engineering and Information Technology* **9** 6 2088-2095
- [11] Situmeang Y P, Suarta M, Irianto I K and Andriani A A S P R 2018 *IOP Conference Series: Materials Science and Engineering* **434** 1 012231
- [12] Situmeang Y P and Suarta M 2018 *Journal of Biological and Chemical Research (JBCR)* **35** 2 463-468
- [13] Situmeang Y P, Sudewa K A and Holo P P 2017 *Journal of Biological and Chemical Research* **34** 2 713-722
- [14] Wirajaya A A N M, Yuliantini M S and Udayana I G B *Proceedings of the 1st Warmadewa Research and Development Seminar (WARDS) European Alliance for Innovation (EAI)*.
- [15] Amaral H D D R, Situmeang Y P, and Suarta M 2019 *Journal of Physics: Conference Series* **1402** 3 033057
- [16] Rahardjo Y C and Purwantari N D 2010 *Potensi Kotoran Kelinci Sebagai Pupuk Organik dan Pemanfaatan Pada Tanaman Pakan dan Sayuran* (Balai Penelitian Ternak)
- [17] Novizan 2005 *Petunjuk Pemupukan yang Efektif* (Jakarta: Agro Media Pustaka)
- [18] Pardosi 2014 Respon Tanaman Sawi Pupuk Cair Limbah Sayuran Pada Lahan Kering Utisol *Prosiding Seminar Nasional Lahan Suboptimal*. Palembang 26-27 September 2014
- [19] Melo L C A, Coscione A R, Abreu C A, Puga A P and Camargo O A 2013 *BioResources* **8** 4 4992-5004
- [20] Situmeang Y P 2018 *MATEC Web of Conferences* **197** 13001
- [21] Brodowski S, John B, Flessa H, and Amelung W 2006 *European Journal of Soil Science* **57** 4 539-546
- [22] Liang B, Lehmann J, Solomon D, Sohi S, Thies J E, Skjemstad J O, Luizão F J, Engelhard M H, Neves E G and Worrick S 2008 *Geochimica et Cosmochimica Acta* **72** 24 6069-6078

Utilization of rabbit manure and biochar chicken manure and its effect on the growth and yield of pakchoy plants

ORIGINALITY REPORT

25%
SIMILARITY INDEX

%
INTERNET SOURCES

25%
PUBLICATIONS

%
STUDENT PAPERS

PRIMARY SOURCES

- 1** U Dani, A N S Budiarti, A A Wijaya. "Application of Chicken manure Dosage and Plant Growth Promoting Rhizobacteria on the Growth and Yield of Shallot Plants (*Allium ascalonicum* L.)", IOP Conference Series: Earth and Environmental Science, 2021
Publication **3%**
 - 2** I G M Arjana, I K A Sudewa. "Study of artificial irradiation arrangement and pinching age on chrysanthemum", IOP Conference Series: Materials Science and Engineering, 2021
Publication **2%**
 - 3** Yedija Manullang, Reny Herawati, Merakati Handajaningsih, Dwi Wahyuni Ganefianti, Eka J.V. Haquarsum, Mimi Sutrawati. "Growth and Yield of Lettuce (*Lactuca sativa* L.) by Giving Cow Manure and Empty Palm Oil Bunch Fertilizer on Peat Soil", Akta Agrosia, 2020
Publication **2%**
-

4

Bo Liang, Shuang Liu, Lina Guo, Dejun Chen, Yong Liu, Zhiyong Zhong, Liufeng Xiong. "Study of humidity on the structure and optical properties of cesium iodide thin film", Journal of Physics: Conference Series, 2016

Publication

2%

5

Herry Gusmara, Ricci Handoko Silitonga, Bilman Wilman Simanihuruk. "Growth Response and Yield of Sweet Corn on Palm Oil Sludge and Dolomite in Ultisols", TERRA : Journal of Land Restoration, 2020

Publication

2%

6

M Khubna, Supriyono, S Nyoto, M T S Budiastuti. "The growth and yield of hybrid corn on different plant spacing", IOP Conference Series: Earth and Environmental Science, 2021

Publication

2%

7

Ignasia Margi Wahyuni, Puspita Ratna Susilawati. "The Use of Teak Leaves (Tectonagrandis) as Alternative Plantation Media for White Oyster Mushroom (Pleurotusostreatus)", Journal of Physics: Conference Series, 2019

Publication

1%

8

M Ansar, Bahrudin, Fathurrahman, S Darman, A R Thaha, A W Angka, Rahmadanih. "Application of bokashi goat manure and

1%

organic liquid fertilizer to improve the growth and yield of Lembah Palu shallot variety", IOP Conference Series: Earth and Environmental Science, 2021

Publication

9

Ryzkita Prima Pramanda, Kuswanta Futas Hidayat, Sunyoto Sunyoto, Muhammad Kamal. "PENGARUH APLIKASI BAHAN ORGANIK TERHADAP PERTUMBUHAN DAN HASIL BEBERAPA VARIETAS SORGUM (*Sorghum bicolor* [L] Moench)", Jurnal Agrotek Tropika, 2015

Publication

10

Hari Soeseno Hardjoloekito. "DOSAGE EFFECTIVENESS OF MANURE AND PLANTING DISTANCE ON THE GROWTH AND PRODUCTION OF PEANUTS (*ARACHIS HYPOGAEA*, L)", International Journal of Research -GRANTHAALAYAH, 2020

Publication

11

Afrah A. Maktoof, Roaa Jafar Elherarlla, Saleem Ethaib. "Identifying the nutritional composition of fish waste, bones, scales, and fins", IOP Conference Series: Materials Science and Engineering, 2020

Publication

12

D Sihombing, D Setyorini, Z Arifin, C Tafakresnanto, W Handayati. "Influence of

1 %

1 %

1 %

1 %

varieties, plant spacing and manure fertilizer to maize productivity in dry land", IOP Conference Series: Earth and Environmental Science, 2021

Publication

13

M. A. K. Khalil. "Heritability of juvenile characters of white spruce (*Picea glauca* (Moench.) Voss.) in central Newfoundland, Canada", Theoretical and Applied Genetics, 1985

Publication

14

F Wibowo, Armaniar. "Physiological performance of the soybean crosses in salinity stress", IOP Conference Series: Earth and Environmental Science, 2018

Publication

15

Izhar Ali, Quan Zhao, Ke Wu, Saif Ullah et al. "Biochar in Combination with Nitrogen Fertilizer is a Technique: To Enhance Physiological and Morphological Traits of Rice (*Oryza sativa* L.) by Improving Soil Physio-biochemical Properties", Journal of Plant Growth Regulation, 2021

Publication

16

Ni Made Yudiastari, Ni Ketut Ety Suwitari, Luh Suariani, Yohanes Parlindungan Situmeang. "The application of biochar in improving the nutrition quality and production of sorghum

1 %

1 %

1 %

1 %

17

Sihufe, G.A.. "The influence of ripening temperature and sampling site on the lipolysis in Reggianito Argentino cheese", Food Research International, 200712

Publication

<1 %

18

A. Fathy Younis, M. A. Hatata. "Studies on the effects of certain salts on germination, on growth of root, and on metabolism", Plant and Soil, 1971

Publication

<1 %

19

Lisa Septiani, RR Yudhy Harini Bertham, Hesti Pujiwati, Bilman Wilman Simanihuruk. "Growth and Yield of Soybean with Application of Liquid Organic Fertilizer and Arbuscular Mycorrhizal Fungi in Ultisols", TERRA : Journal of Land Restoration, 2021

Publication

<1 %

20

Hafiza Iqra Khalid, Ghulam Murtaza, Saif Ullah, Ejaz Ahmad Waraich. "Nanoparticles Induced Restricted Cadmium Mobility and Accumulation in Wheat From Contaminated Soil", Research Square Platform LLC, 2021

Publication

<1 %

21

A T Maryani, N Mirna, F F Bahar. "The effect of organic fertilizer chicken on the growth of

<1 %

papaya plants in palm oil plantation: a case study of Naga Sari Village, Mestong, Muaro Jambi District", IOP Conference Series: Earth and Environmental Science, 2021

Publication

22

Sri Pratiwi Aritonang, Ernitha Panjaitan, Fetrus Parsaulian Tondang. " Cucumber plants (I.) growth and crop yield of chicken manure fertilized with plant spacing ", IOP Conference Series: Earth and Environmental Science, 2018

Publication

23

I G B Udayana, L Kartini, Y P Situmeang, I K Sunadra. "The modeling system of horticultural commodity development in Pinggan village, Bangli regency, Bali Province", Journal of Physics: Conference Series, 2019

Publication

24

Rezvan Ramezannejad, Morteza Mofid Bojnoordi, Mohammad Armin, Mahnaz Aghdasi. "Artificial Neural Network Modeling To Predict And Optimize Phenolic Acids Production From Callus Culture of Lactuca Undulata", Research Square Platform LLC, 2021

Publication

25

Sugeng Winarso, Bambang Hermiyanto, Sukron Romadhona, Martinus H. Pandutama, Tri Candra Setiawati, Indasah. "Effectiveness

<1 %

<1 %

<1 %

<1 %

of Biopellet Combination of Biochar, Chicken Manure and Fish Waste To Improvement of Chemical Properties of Sandy Soil and Soybean Plant Growth", IOP Conference Series: Earth and Environmental Science, 2021

Publication

26

Ángela Gómez Gómez. "Pasteurization of Lipid Emulsions with Supercritical CO₂ and High Power Ultrasound", Universitat Politecnica de Valencia, 2021

Publication

<1 %

27

Xixi Liu, Xiaoxun Jiang, Weirong Zhao, Yuanyuan Cao, Tingting Guo, Xiangyi He, Haiting Ni, Xinyun Tang. " Colonization of phosphate-solubilizing sp. strain P34-L in the wheat rhizosphere and its effects on wheat growth and the expression of phosphate transporter gene in wheat ", Cold Spring Harbor Laboratory, 2018

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Utilization of rabbit manure and biochar chicken manure and its effect on the growth and yield of pakchoy plants

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
